

THE IN A HURRY SERIES

Deserts *in a Hurry*

Scarcity, survival, and hidden abundance

Max Brocklesby

BOOKSINAHURRY.COM

Contents

The Whole Thing in One Page 4

Why You Should Care 5

The Core Ideas 7

How It Actually Works 20

What People Get Wrong 30

Use It 35

Terms 39

Go Deeper 43

Notes and Sources 44

Bibliography 51

The Whole Thing in One Page

The desert in your head is hot, sandy and empty. None of those words defines one. Antarctica is a desert. Much of the Sahara is rock, gravel, mountain, dry valley or salt depression rather than dune. Ground that appears vacant may hold seeds waiting for rain, animals asleep below the heat, a crust of living organisms, a route used in another season, or groundwater that fell when the climate was wetter.

A desert is a place where usable moisture is persistently scarce. Low rainfall can create that condition, but so can fierce evaporative demand, cold conditions that limit atmospheric moisture, mountains that strip moisture from winds, cold ocean currents that suppress rain, or great distance from the sea. These causes often combine. Heat intensifies some deserts. It does not define them.

Scarcity does not stop the system. It changes its rhythm and geometry. Water arrives in pulses, then gathers in channels, fans, springs, fog belts, snowfields and aquifers. A dry wash can become a lethal river beneath a storm that fell beyond the horizon. A few hours of rain can wake microbes, germinate annuals, fill temporary pools and support a season of breeding. The important contrast is seldom wet against dry. It is a brief event against the long interval that follows.

Life survives by altering the exchange. CAM plants take in carbon dioxide mainly at night and close their pores through the hotter day. Other plants spread shallow roots, reach deep water, shed leaves or wait as seed. Animals burrow, move after sunset, tolerate changing body temperature or suspend activity. A camel's hump contains fat. Its endurance comes from an integrated physiology, joined to people who know where forage and water lie.

The ground is working too. Floods build alluvial fans. Wind moves sand and dust. Evaporation concentrates salts. Biological soil crust binds exposed particles. A tyre track can outlast the journey that made it. What looks blank may be an archive, a habitat or both.

People learned to live with dispersion. Pastoral mobility joined seasonal pasture to wells and fallback ground. Caravans joined water points to markets. Qanats carried groundwater downhill through tunnels, but labour, rules and repair kept the flow alive. Modern cities stretch the same problem across dams, pipelines, power grids, treatment works and legal claims.

This is the hidden abundance in the subtitle: value concentrated in rare places, brief seasons and old stores. Springs can hold species found nowhere else. Closed basins collect minerals. Clear skies favour solar power. Deep aquifers support farms and cities. Concentration makes each resource useful and each loss consequential.

The final danger follows from the first condition. A system adapted to slow renewal can look generous while its reserve is being spent. Pump an aquifer faster than recharge, salt irrigated soil, block a grazing corridor, drain a spring or build across intact habitat because it looks unused, and scarcity becomes depletion. Desertification is neither the natural existence of desert nor one wall of sand advancing. It is lost land function in drylands through climatic pressure and human activity.

To understand a desert, track why moisture is scarce, when it arrives, where it gathers, who or what can reach it, and how long renewal takes. The emptiness then becomes a map of pulses, stores, refuges and routes.

That is the book.

Why You Should Care

On 20 August 2023, the remnants of Hurricane Hilary crossed Death Valley. The park normally receives little rain across a whole year. That storm delivered 2.2 inches at Furnace Creek, tore roads apart and sent mud and stone through channels that visitors had seen only as dust. Water spread across Badwater Basin, the lowest ground in North America. Further rain in February 2024 refreshed it. For several weeks, people paddled on a temporary lake reported at roughly six miles long, three miles wide and one foot deep.

The lake, informally called Lake Manly after the ancient lake that once occupied the basin, was not a contradiction. It was a desert event. Scarce water arrived as destructive excess, revealed the shape of the catchment, settled in the lowest hollow and then thinned under sun and wind. Important processes in deserts are often separated by long quiet intervals. When they occur, they can be faster and larger than the empty scenery prepared you to expect.

This matters because modern life is good at mistaking a stable output for a stable system. A desert farm can harvest for decades while its water table falls. A city can look detached from scarcity because snowmelt, electricity and food arrive from far away. A rangeland can seem unused during one season while remaining essential after the next failed rain. An annual average can be correct and still conceal the sequence that decides survival.

Deserts also expose the cost of bad categories. Calling every dry spell desertification confuses weather, climate and damage. Calling mobile pastoralism wandering turns a structured response to variable pasture into a planning defect. Calling a spring small because it occupies little area misses the species, herds and households concentrated there. Calling land empty makes residents and seasonal users disappear before any bulldozer arrives.

A 2024 assessment by the United Nations Convention to Combat Desertification compared the climate of 1991 to 2020 with 1961 to 1990. It found that the area classed as dryland had grown by about 4.3 million square kilometres to cover more than 40 per cent of land excluding Antarctica. Drylands include semi-arid grasslands, farms and settlements as well as deserts, so this is not a claim that two-fifths of the planet is sand or ruined. It is a measure of long-term atmospheric aridity, not a direct map of degradation. The distinction matters because climate can tighten the margin while land rights, farming, grazing, extraction and infrastructure decide what happens next.

The subject also reaches beyond dry-country borders. Dust from North Africa crosses the Atlantic and carries nutrients to distant ecosystems.

Desert basins contain copper, lithium brines, salts, oil and gas. High solar exposure places some regions at the centre of energy plans. Rivers born in wet mountains sustain cities and farms far downstream in desert climates. A place coloured pale on a vegetation map may sit at the centre of a supply chain.

The lesson is not that every desert should be left untouched or every old practice preserved. Irrigation can support flourishing societies. Desalination can secure coastal water. Solar fields can displace fossil generation. Grazing can maintain livelihoods or damage land, depending on timing, access and pressure. The useful discipline is to identify the whole system before judging the visible patch, then distinguish the resource that returns from the reserve that does not.

By the end of this hour, you should be able to look past dunes and heat. You should see the air that withheld rain, the channel waiting for a storm, the seed waiting for its cue, the well connected to a distant recharge zone, the route joining sparse resources, and the old store whose apparent plenty depends on a forgotten climate. A blank part of the map will no longer look blank.

The Core Ideas

A Desert Is a Moisture Balance, Not a Temperature

Rainfall is the wrong place to stop. A landscape receiving 200 millimetres of rain beneath cool cloud does not face the same water problem as one receiving the same amount under fierce sun, hot air and persistent wind. What matters is the balance between moisture supplied and the water that the atmosphere, soil, plants and streams can remove. The shared condition is a persistent deficit.

This is why the familiar threshold of about 250 millimetres of annual precipitation is useful but crude. It gives a rough map, yet it misses fog, snow, seasonality, runoff and evaporative demand. Geographers therefore

use measures such as the aridity index, which compares precipitation with potential evapotranspiration. It asks how much moisture arrives relative to how much the atmosphere could remove if water were available. Even that ratio cannot show every spring, channel or season. A shaded canyon can remain moist inside a dry region. A mountain storm can recharge a basin that receives little rain on its floor. Classification draws a climatic boundary; organisms and households live inside the local pathways.

Temperature is secondary. Antarctica receives so little precipitation that its interior is a polar desert despite being covered by ice. The Sahara is the largest hot desert, but hot is a modifier, not the definition. The Gobi has severe winters. High plateaux in Central Asia and the Andes combine cold, wind, altitude and dryness. A desert can freeze a person or cook one. Some can do both within a day.

The definition also separates desert from drought. A desert is a long-term climatic and ecological condition. Drought is an interval when water falls below what a place, population or system normally expects. A desert can suffer drought, and a wet region can suffer one too. The same reservoir level may be ordinary in one basin and an emergency in another because the baseline, demand and alternatives differ.

Dryland is broader again. In broad geographical classification, drylands span hyper-arid, arid, semi-arid and dry sub-humid climates, though the United Nations desertification convention excludes hyper-arid and polar regions from its operational definition. Drylands contain deserts, grasslands, farms and towns. Desertification concerns degradation in defined dryland zones rather than the appearance of classic desert scenery. Aridification means a long-term shift towards a drier climate. These categories answer different questions: what the climate is, whether it has changed, whether land has lost function, and whether current supply has departed from normal.

Once desert is treated as a moisture balance, several puzzles clear. A place can flood and remain a desert because a violent event does not erase a long-term deficit. A coast can be foggy while receiving almost no rain. A city

can look green while pumping water accumulated under an older climate. The controlling questions are how water enters, where it travels, how long it stays and which lives or uses can reach it. Those questions define the rest of the book.

There Is More Than One Way to Lose the Rain

The great hot deserts cluster near 20 to 30 degrees north and south because the atmosphere has circulation. Warm, moist air rises near the equator, cools and releases much of its water. Higher in the atmosphere it moves poleward, then descends in the subtropics. Descending air warms as pressure increases. Its relative humidity falls, clouds struggle to form and rain is suppressed. The Sahara, Arabian Desert and much of Australia lie beneath these broad belts of subsiding air.

That is one engine, not the engine. Mountains create another. Air forced up a windward slope cools, condenses and rains. By the time it crosses the crest, it has lost moisture. Descending on the leeward side, it warms and dries. The result is a rain shadow. Patagonia lies east of the Andes. The Great Basin lies beyond the Sierra Nevada. Parts of Central Asia sit behind several mountain barriers, each taking another cut from the air before it arrives.

Cold ocean currents help create a stranger desert. Along the western edges of continents, cold water cools the air immediately above it. A stable inversion can trap that cool layer beneath warmer air, discouraging the vertical growth of rain clouds. The Namib and Atacama may receive fog while remaining among the driest places on land. Fog is not failed weather there. It is a separate water supply, harvested by beetles, plants and, increasingly, human collectors. A landscape can be wet to the touch and remain arid over the year. The Atacama shows how mechanisms can stack: cold Pacific water, stable descending air, mountain barriers and great distance from Atlantic moisture combine to suppress rain with unusual force.

Continental interiors lose rain through distance. Air masses travelling far from oceans shed moisture on the way. The Gobi and Taklamakan also sit

behind mountains, so continentality and rain shadow reinforce each other. Polar deserts reach dryness by cold. At polar temperatures, the atmosphere reaches saturation with little water vapour, limiting snowfall even above a vast ice sheet. High altitude adds another route by cooling the atmosphere and shortening the growing season.

These causes combine and shift. The Sahara has not always looked as it does. Changes in Earth's orbit alter the distribution of sunlight, which changes monsoons over thousands of years. During the African Humid Period, lakes, rivers, grasslands and people occupied regions now hyper-arid. Rock art records cattle and wildlife where water is now scarce. The return of aridity was not a single morning when grass became sand. Rain belts moved, lakes shrank, vegetation changed and people reorganised or left over generations.

This historical movement matters because it prevents a false choice between natural desert and human damage. Deserts are genuine climatic regions, not mistakes that people made. Human action can still degrade dryland, change dust emissions, drain wetlands or exhaust groundwater. A natural origin does not make every present condition natural, and a damaged dryland does not prove that all aridity is damage.

The same landscape may therefore owe its dryness to global circulation, mountains, ocean temperature, distance from moisture, altitude and past climate at once. Any explanation that assigns one cause to all deserts will fail somewhere. The shared result is scarcity. The route to it remains local.

Water Arrives in Pulses and Persists in Stores

In many deserts, the important unit of rainfall is not the year. It is the event. A long dry interval may be broken by a short storm intense enough to exceed the ground's ability to absorb water. Bare slopes, crusted soils and steep channels send runoff into wadis, arroyos and washes. A channel that carried dust in the morning can carry boulders by afternoon. People drown in desert floods because dryness encourages the belief that the channel is a road. The damage depends on more than the storm total. Rainfall intensity, slope, soil condition, channel shape and buildings placed in the flow path decide whether water infiltrates, spreads or strikes as a moving load of sediment.

The same flood can create the next reserve. Water spreads across an alluvial fan, slows and sinks through gravel. It fills shallow sediment, reaches fractures in rock or feeds groundwater at the edge of a basin. Recharge is patchy. A gauge on open ground may record little useful infiltration while a channel several kilometres away supplies an aquifer. Desert hydrology is concentrated by terrain.

Ecologists describe a related pattern as pulse and reserve. Rain triggers germination, photosynthesis, microbial activity, insect emergence and breeding. Organisms turn the temporary surplus into stored seed, tissue, fat, eggs or nutrients. Activity then contracts as the surface dries. The reserve carries the system towards the next pulse. This is not a machine that switches on uniformly. A five-millimetre shower may wet the surface and feed microbes but fail to reach deep roots. A larger storm may penetrate further and sustain shrubs for months. Timing, depth and sequence matter as much as the annual total.

Water also hides in forms that confuse the eye. It may sit beneath an alluvial plain, gather above impermeable rock, emerge at a fault as a spring, arrive as fog on a coastal slope, or remain frozen in a cold desert. Some deep aquifers contain water that fell during wetter climatic periods. Calling all such water fossil is imprecise, but the management question is plain: if modern recharge is negligible compared with pumping, the reserve behaves like an

inherited store rather than an annual supply.

An oasis forms where hidden water becomes accessible, but the green patch is rarely self-explanatory. Springs can feed one. A high water table can support wells. Floodwater can be diverted. Qanats can intercept an aquifer upslope and carry water underground. Soil, labour, ownership and maintenance decide how much greenery follows. A cultivated oasis is a water institution with trees around it.

Evaporation concentrates what water leaves behind. Closed basins have no river outlet to the sea. Water enters, dissolves minerals, then disappears into air. Salts remain. Repetition produces playas, salars and salt crusts, along with deposits of borates, potash and lithium-bearing brines in some basins. Hidden abundance is produced by loss.

The model changes how scarcity should be measured. Counting wells without measuring water levels says little. Recording rainfall without mapping channels misses recharge. Taking a long-term average can erase the few events on which the system depends. Desert water must be followed through time and through the ground. What vanishes from sight may have been lost, stored or merely moved.

Life Survives by Changing the Clock

A cactus cannot outrun noon, but it can alter when it trades with the air. Most plants take in carbon dioxide through pores called stomata. Open pores also lose water. Many succulents use crassulacean acid metabolism, or CAM, opening stomata at night when air is cooler and often more humid. They store the captured carbon in organic acids, close the pores by day and use the stored carbon for photosynthesis. The system saves water, but it limits the rate at which carbon can be acquired. Survival is bought with slower growth.

Other plants choose different bargains. Some spread shallow roots far beyond the visible crown to seize brief showers. Others send roots deep towards more reliable moisture. Tiny leaves, waxy surfaces, hairs and reflective coatings reduce heat load or water loss. Drought-deciduous

shrubs shed leaves when soil moisture falls. Annual plants avoid the dry season almost entirely. The adult dies, while seeds wait in soil until rainfall, temperature and season align. After a strong event, a plain that looked empty can flower because the population was present in another state.

No strategy is universally best. Succulence stores water but creates valuable tissue for animals and can fail under freezing conditions. Deep roots cost carbon to build. Shallow roots capture light rain but compete intensely near the surface. Dormancy avoids bad years but risks missing the rare good one if germination cues are wrong. Desert evolution does not produce perfection. It produces workable compromises under recurring shortage.

Animals can leave the heat without leaving the desert. Burrows buffer temperature and humidity. Nocturnal and crepuscular activity shifts foraging away from midday. Small mammals may obtain much of their water from food and metabolic processes while producing concentrated urine and dry faeces. Reptiles allow body temperature to vary and move among sun, shade and shelter. Some amphibians remain dormant underground through long dry periods, emerging after rain to feed and breed at speed. Heat and water are coupled: evaporative cooling spends water, so an animal that avoids heat can conserve moisture without changing its kidneys. Behaviour is physiology performed with movement.

Large animals face different physics. A camel's hump stores fat, not free water. The concentrated reserve supplies energy when forage is poor without turning the hump into a water cistern. Camels also tolerate substantial dehydration, reduce urinary and faecal water loss, reclaim moisture in nasal passages and allow body temperature to vary, delaying the need to sweat. No single trick explains the animal. The myth survives because a tank is easier to remember than an integrated physiology.

The most dramatic response to a pulse is often reproduction. Temporary pools can fill with invertebrates whose eggs survived in dry sediment. Insects emerge, birds arrive, frogs call and predators follow. The boom may be brief enough that an observer visiting in another month sees none of it.

Desert biodiversity is easily underestimated when surveys are timed to human calendars rather than ecological ones.

This changing clock explains why disturbance can be severe even where recovery appears possible. A plant may grow slowly for decades. A soil crust may take years to rebuild lost structure. A tortoise population may occupy a wide area at low density. The desert can produce spectacular bursts and still repair physical damage slowly. Pulse does not mean quick replacement of everything.

Scarcity therefore shapes life through timing as much as anatomy. The successful organism is not always the one that endures the harshest moment in full activity. Often it is the one that knows, through physiology rather than thought, when to close, hide, wait or begin.

The Surface Is Alive and Mostly Not Sand

Dunes own the photographs because wind gives them clean lines. They do not own most desert ground. Many deserts are dominated by bare rock, gravel plains, alluvial fans, dry channels, clay pans, salt flats and low shrubland. Even within the Sahara, sand seas are one part of a much larger mosaic. Sand is mobile and visible. Stone and crust are quieter, so popular memory mistakes visibility for extent.

Water builds much of this supposedly wind-made world. Rare floods cut channels, move coarse sediment out of mountains and spread it in fans where slopes flatten. Repeated flows join fans into broad aprons. In closed basins, fine sediment settles in playas. When water evaporates, salts crystallise. Wind later lifts the finest exposed particles as dust or moves sand grain by grain. The landscape is an argument between rare water and persistent air. Temperature and salt work more slowly. Repeated heating, cooling, wetting and crystallisation weaken exposed rock. With little vegetation to hide the result, cliffs retreat, blocks break and old surfaces remain legible. Desert scenery can look freshly stripped while containing landforms assembled over immense spans of time.

A dune needs three things: sand, wind capable of moving it and a place

where transport loses efficiency. Its shape records wind direction, sand supply and vegetation. Barchans form crescent shapes under mainly one wind direction and limited sand. Linear dunes record a more complex wind regime. Star dunes grow where winds arrive from several directions. Dunes migrate, but not every sand sea is racing across the map. Some are anchored by plants, moisture or a stable wind balance.

Gravel surfaces can be ancient. Desert pavement consists of closely packed stones above finer material. Several processes contribute to its formation, including the trapping and downward movement of dust and the reorganisation of particles near the surface. Once established, the pavement shields soil from wind. A vehicle track can break that armour and expose fine sediment. The mark may persist far longer than the journey that made it.

Then there is the ground that looks least alive. Biological soil crusts are communities of cyanobacteria, algae, fungi, lichens and mosses living on or just beneath the surface. Filaments bind particles. Some organisms fix carbon or nitrogen. Crusts alter infiltration, retain material and reduce erosion. Their effects vary with species, soil and climate, so they are not a universal magic skin. They are still evidence that the apparent gap between shrubs can be occupied habitat.

Dust gives the surface a global reach. Fine mineral particles lifted from dry lake beds and disturbed soils can travel across continents and oceans. They affect air quality, clouds, snowmelt and nutrient cycles. Using satellite observations from 2007 to 2013, researchers estimated that African dust carried about 22,000 tonnes of phosphorus to the Amazon basin per year, with a wide uncertainty range. They judged the input comparable to estimated hydrological losses of phosphorus. The number is neither constant nor a complete account of Amazon fertility. The firmer lesson is that material exported from dry land can move nutrients between distant ecosystems.

Desert surfaces are therefore archives as well as habitats. Old shorelines mark vanished lakes. Fans record floods. Salt crusts record evaporation.

Pavements record stability and dust. Tracks record disturbance. The blankness is partly a failure of vocabulary. Once the surfaces are named, they stop looking interchangeable.

Human Life Depends on Routes, Rights and Maintenance

People did not wait for modern engineering to enter deserts. Archaeology and living practice record hunting, farming, pastoralism, settlement and trade across dry regions that outsiders later labelled empty. The error comes from looking only for the permanent marks of settled agriculture. A field leaves straight lines. Mobility leaves knowledge, agreements, remembered water points and seasonal presence, much of it harder to see from a surveyor's desk. A route may be infrastructure even when nobody poured concrete.

Local knowledge turns sparse ground into a detailed map. Gary Paul Nabhan's account of O'dham country describes Tohono O'dham farming and gathering practices that match scarce water, cultivated crops and wild foods to particular places and seasons. The example shows how land described as inhospitable by newcomers can be densely legible to people who know its storms, plants and intervals. It is neither a universal desert method nor proof that inherited practice cannot fail. It shows that scarcity is partly a problem of access, memory and maintenance, not rainfall alone.

Pastoral movement is another response to variability. Rain may produce good grazing in one district and miss the next. Wells fail at different times. Herds need to reach forage without exhausting one patch. Routes connect resources that are individually inadequate but collectively usable. Movement can spread pressure and allow recovery, yet it can also contribute to conflict or degradation where access is blocked, stocking is excessive or drought is prolonged. The important correction is that mobility has structure. It is not the absence of management.

The camel enlarged what those networks could carry. Its physiology allowed longer intervals between water, while its feet and load-bearing capacity

suited sandy and stony ground. Caravan trade still required wells, grazing, guides, protection, credit and staging points. Across the Sahara, salt, gold, cloth, books and enslaved people moved between West Africa, North Africa and Mediterranean markets. Timbuktu grew near the southern edge of those routes because water, political authority and exchange met there. The desert did not disappear. Distance became organised.

Permanent settlement depended on networks too. The Persian qanat begins with a mother well reaching groundwater at higher ground. A gently sloping tunnel carries water towards fields and settlements by gravity, with vertical shafts providing access for construction and repair. Keeping the channel underground reduces evaporation and contamination. The tunnel does not create water. Its success depends on the water table, continual labour, allocation rules and restraint at the source.

Yazd shows the larger fit. Thick earthen walls, shaded alleys, courtyards, cisterns, windcatchers and qanats reduced particular losses of heat and water. The city was not independent of the desert. It was closely adjusted to it.

Scarcity is also divided inside households. In countries with individual-level data, women and girls were responsible for water collection in seven out of ten households where water had to be fetched away from home. That work can vanish from a map of wells or pipes even though distance is paid in time and physical effort. Water access is a question of labour and power as well as litres.

Many modern desert cities pursue a different strategy by extending their supply network outward. Rivers are dammed far away. Groundwater is pumped, food imported and electricity used for cooling or desalination. Wastewater may be treated and returned. In southern Nevada, treated indoor water sent back to Lake Mead earns return-flow credit, while much outdoor irrigation is lost through evaporation and plant transpiration. This is a local arrangement created by topography, treatment works and Colorado River law, not a general rule for desert cities.

The human desert is therefore best mapped through water points, pasture, tunnels, routes, pipes, markets, labour and rights. A gap between nodes may be the space that keeps movement possible. Cut a corridor, monopolise a well or neglect a channel, and a low-density system can fail without any single place looking crowded. Survival is social before it is picturesque.

Hidden Abundance Can Be Spent Faster Than It Returns

Deserts concentrate valuable things because water and vegetation do less to cover, dilute or carry them away. Evaporation leaves salts. Ancient lakes and hydrothermal systems contribute mineral deposits. Clear skies and open ground favour solar generation. Sedimentary basins may hold oil and gas. Deep aquifers can support cities and farms. Concentration can be biological too: an isolated spring may hold species found nowhere else, while a salt flat used by industry may belong to a wider wetland system. Value and vulnerability gather in the same places.

Groundwater is the clearest case of apparent plenty. A 2024 analysis of 170,000 monitoring wells in 1,693 aquifer systems found rapid groundwater-level declines widespread in the twenty-first century, especially in dry regions with extensive cropland. The data came from more than 40 countries that together account for about three-quarters of global groundwater withdrawals, but the wells were not a random global sample. The study also found cases where policy change, managed aquifer recharge or surface-water diversions slowed or reversed declines. The lesson is neither universal collapse nor easy recovery. Local wells and aquifer boundaries remain essential because nearby trends can differ and satellite gravity is too coarse for local management.

Irrigation creates another delayed failure. Water applied to crops evaporates or is transpired, leaving dissolved salts behind. If drainage is poor, salts accumulate around roots and yields fall. More water may flush them downward, but that requires supply and somewhere for saline drainage to go. Irrigation can sustain desert agriculture for generations. Green fields

alone cannot show whether the soil and water system is renewing.

Mineral extraction concentrates benefits and costs unevenly. Lithium production in high Andean closed basins uses brine and, in varying amounts, freshwater. A 2025 study estimated modern freshwater inflows for 28 active or prospective lithium-producing basins in Chile, Argentina and Bolivia at 2 to 33 millimetres a year. It found that common global hydrological models substantially overestimated freshwater availability in this setting. Freshwater pumping, brine pumping and process-water use are related but not interchangeable measures; their effects can differ with hydrogeology and distance from wetlands. Direct streamflow measurements were available for only three basins, so the result is not a universal impact number. It demonstrates the need to assess freshwater use, brine abstraction, wetlands, community supply and the position of pumping points basin by basin.

Solar energy presents a related trade-off. Strong sunlight makes many deserts attractive for low-carbon power, yet utility-scale facilities occupy land, build roads, fragment habitat, disturb soil and alter drainage. Conservation planning in the Mojave shows why siting matters, but its species maps and mitigation options cannot be copied unchanged elsewhere. Rooftops, degraded land, transmission access and habitat differ. Empty-looking ground is not impact-free ground.

Desertification is the broadest failure. Under the United Nations convention, the term covers lost land function within arid, semi-arid and dry sub-humid zones where climatic variation and human activity contribute. It includes reduced biological or economic productivity and ecological complexity. It does not require dunes. Overgrazing can contribute in one place, while boundaries that prevent adaptive movement can concentrate grazing in another. Irrigation salinity, tree removal, repeated burning, mining, groundwater decline and warming may interact. There is no single desertification machine.

Aridification adds pressure without proving degradation. A 2024 United Nations assessment compared consecutive thirty-year climate periods and

found a marked expansion in the area classified as dryland by 1991 to 2020. The result describes long-term atmospheric aridity, not one advancing front or a direct inventory of damaged ground. Regional causes and consequences differ.

Restoration works when it repairs processes rather than paints the surface green. That may mean protecting biocrust, slowing runoff, improving drainage, securing pastoral routes, changing stocking rules, reducing groundwater withdrawal or allowing passive recovery. Planting trees where water cannot support them can deepen scarcity. A sound intervention asks what function was lost, at what scale and according to whose evidence.

The causal loop is complete. Persistent scarcity concentrated water, life and human activity into stores, nodes and routes. Those concentrations made desert wealth usable. Modern pumps, roads and large capital projects can enlarge access and withdrawal until a slow reserve behaves like a temporary flow. The tools that overcome scarcity can therefore expose the system to depletion. Renewal time is the limit that technology can postpone but cannot remove.

How It Actually Works

Before a drop falls

A desert water cycle begins with air that fails to deliver. Across the subtropics, descending air warms, lowers relative humidity and suppresses cloud. Behind a mountain, air has already surrendered much of its moisture on the far slope. Beside a cold current, a shallow layer of damp air may sit beneath an inversion, producing fog without the rising motion needed for rain. In a polar desert, cold air reaches saturation with little water vapour, so snowfall remains scant.

The absence is never complete. Moisture still arrives through storms, snow, fog, dew and rivers born outside the desert. The Nile crosses the Sahara because its headwaters are elsewhere. The Colorado reaches dry country with water gathered from mountain snow. Coastal fog wets plants in the

Namib and Atacama. Each source enters on a different schedule and answers to a different climate. Treating them as one supply called water hides the engineering problem.

Topography divides the incoming moisture before any person can. Ridges force air upward. Basins collect runoff. Bedrock sends water sideways along fractures. Gravel stores it. Clay slows it. Salt changes what can live in it. Two valleys receiving the same storm may finish with different floods, different recharge and different vegetation because the pathways beneath the rain are different.

Season adds another division. The Sonoran Desert receives winter storms from the Pacific and summer monsoon rain from the south, giving organisms two possible growing periods. Mediterranean-climate deserts receive most of their rain in cooler months. Polar deserts may store snowfall for years before meltwater moves. A yearly total cannot show whether water arrived as gentle winter rain, a convective downpour, fog droplets or snow. Each enters the system through a different door.

Rivers born beyond the climatic boundary complicate the map further. The Nile, Indus and Colorado carry imported water through dry regions, creating corridors whose productivity reflects remote mountains and rainfall. The river valley may be lush while the surrounding basin remains desert. Political borders usually follow neither catchments nor aquifers, so the physical source and the institution claiming it may be far apart.

A storm meets the ground

The first drops may vanish into hot, dry air or dampen only the top millimetres of soil. If rain continues, the surface begins to decide. Loose sand can absorb water rapidly, though water may not stay near roots. Fine soil can seal as raindrops strike it. Biological crust can increase or reduce infiltration depending on its composition and setting. Rock absorbs little. A steep slope turns excess into speed.

Once rainfall exceeds infiltration, water runs. Small threads join, channels fill and the flood front advances into ground where no rain may have fallen. In

arid basins this can happen with little warning. Runoff carries sand, gravel, branches and debris, so the hazard is not a clean rise in water level. It is a dense moving mixture capable of cutting roads and shifting channels.

At the mouth of a mountain canyon, the slope eases. The stream loses carrying power and drops its coarsest load. Repeated floods build a fan-shaped deposit. Channels split and migrate across it. Houses placed on an apparently inactive part of the fan may stand on the path selected by a later flood. The long dry interval makes old channels look safer than they are.

Some water spreads into a closed basin. A shallow lake may form, sometimes for days and sometimes for years. Without an outlet to the sea, the basin loses water mainly through evaporation or infiltration. Mud settles. Dissolved minerals remain. When the surface dries, it becomes a playa or salt flat. Wind can then lift exposed fine sediment, linking the flood cycle to the dust cycle.

People have long used the violence selectively. Spate irrigation diverts short-lived floodwater onto fields, where sediment and moisture are spread before the channel dries. Small barriers can slow runoff and encourage infiltration. Both methods require attention to flood size. A structure that captures a modest flow may fail under a rare large one or deprive users downstream. Controlling every pulse can also starve wetlands, fans and aquifers that depend on occasional overflow.

The sequence explains why a rain gauge cannot stand in for flood risk. A storm over a mountain may send water towards a town under clear sky. The channel remembers the whole catchment. The person standing in it sees only the patch overhead.

The green switch

Rainfall begins several clocks. Cyanobacteria and other organisms in the soil surface become metabolically active. Seeds take up water. Annual plants germinate if temperature and season provide the right cue. Perennial shrubs extend leaves or roots. Insects emerge from eggs or dormant stages. Temporary pools fill with organisms whose life cycles are built for urgency.

The response is uneven because water penetrates to different depths. A light shower may activate microbes and shallow-rooted plants, then disappear within days. A larger pulse reaches perennial roots and supports growth for longer. Closely spaced storms can produce more biological effect than the same total scattered across months, because the second event arrives before the first has been lost. Conversely, a warm spell after rain can remove the gain rapidly.

Plants turn the pulse into reserve. An annual makes seed. A shrub adds tissue or stores carbohydrates. A succulent replenishes water. Animals convert food into fat, eggs or young. Predators follow prey. Migrating birds use wetlands and blooms that exist only briefly. The desert appears to erupt, but the event is the visible phase of a system that spent most of its time prepared.

Failure is part of the same mechanism. Seeds can germinate after a false start and die before reproducing. A storm may arrive too late for pollinators or too early for warm temperatures. A flood can scour plants from a channel. Desert life is adapted to variability at population scale, not protected from every bad sequence. Long waiting and sudden loss coexist.

The food web follows the geometry of moisture. Productive strips form along washes, beneath shrubs and around springs. A shrub catches windblown litter, shades soil and concentrates nutrients, creating an island that supports insects and smaller plants. A wetland no larger than a few fields can become the only reliable water for kilometres. Predators learn the same map as prey. The ecosystem is sparse in average cover but dense in

consequence at its nodes.

This patchiness affects monitoring. A survey line placed between shrubs can report low productivity while missing the channels that carry most growth. A visit after a failed rainy season can miss species present as seed, eggs or dormant adults. Repeated observation across events is necessary because presence and activity are not the same thing.

Into the ground

Water that infiltrates enters a second landscape. Pores between grains hold it against gravity near the surface. Below that zone, water moves downward or sideways according to permeability and pressure. Gravel and fractured rock can transmit it. Clay and unfractured rock impede it. Where an aquifer meets the surface, a spring or wetland may form.

Recharge often occurs far from the well that later withdraws the water. Mountain fronts receive more precipitation than basin floors. Floods in ephemeral channels can leak through coarse sediment. Snowmelt can enter fractures. Isotopes and dissolved gases sometimes show that deep water fell under climatic conditions unlike those of the present. The age of water is not the same as the age of the rock holding it, and one sample cannot describe a whole aquifer.

An oasis appears when people, plants or animals can reach this hidden store. Springs are obvious points. Shallow groundwater can be lifted from wells. Date palms can create shade above fruit trees and crops, but the familiar layered garden depends on soil, salt control and labour as much as water. Irrigation schedules and ownership rules decide who receives which share. Maintenance decides whether channels silt up. The green patch is the surface expression of a managed flow.

Salinity accumulates as water cycles through. Irrigation water contains dissolved minerals even when it tastes fresh. Plants use water and leave much of the salt. Evaporation does the same. Drainage must carry salts below roots or out of the field. In a closed basin there may be no easy destination. A productive irrigation system therefore needs drainage as

much as delivery.

Groundwater links surface users who may never meet. Pumping near a spring can reduce flow kilometres away. A deep well can lower pressure in shallower layers if the formations connect. Conversely, an aquifer may be compartmentalised by faults or clay, so decline in one part does not describe the whole basin. The word aquifer encourages the image of an underground lake. Most are water held within pores and fractures, with boundaries that must be inferred.

Springs reveal the ecological cost of that uncertainty. Fish, snails, plants and insects may have evolved in isolated waters with nowhere else to go. A small reduction in discharge can change temperature, chemistry and habitat before the spring disappears. The amount withdrawn can look minor at regional scale and decisive at the outlet.

From water point to route

A single well cannot support a mobile livelihood unless it connects to pasture, other wells and rights of passage. Pastoral routes are built around seasonal expectation and contingency. A herd may move towards areas that received rain, avoid disease, reach market or preserve dry-season grazing. Decisions combine observation, messages from other herders, customary agreements and memory of previous years.

Movement changes the pressure on land. Used well, it allows livestock to follow dispersed forage and gives grazed patches time to recover. Used under constraint, it can concentrate animals around permanent water, roads or fenced boundaries. The same number of animals can have different effects depending on where and when they remain. Counting livestock without mapping movement is another desert average that conceals the mechanism.

Trade routes enlarged the network. A caravan required animals, fodder, water, guides, protection, finance and knowledge of intervals between stops. The distance between wells set load and timing. Salt moved south across the Sahara while gold and other goods moved north, alongside

books, cloth and enslaved people. Towns grew where routes, water and political authority met. Their location was a solution to connection, not a refusal of geography.

Qanats converted an aquifer into a linear settlement system. Workers sank a mother well to reach groundwater near higher ground, then excavated a gently descending tunnel towards fields and houses. Vertical shafts removed spoil and allowed repairs. Gravity supplied the energy. The underground channel reduced evaporation, but it did not create water. If the water table fell below the tunnel or maintenance failed, the flow declined. The technology succeeded by fitting extraction to slope and collective upkeep.

Information moved through these networks with goods. Guides knew the reliability of wells, the condition of pasture, the politics of a crossing and the pace of different animals. Market news changed what a caravan carried. Kinship and commercial ties provided credit and protection. Calling a route a line on a map strips away the institutions that made the line usable.

The same is true of mobile pastoralism. Herders often keep several options rather than optimise one fixed circuit. A reserve pasture may be left unused in ordinary years because its value appears only during drought. Formal land allocation can erase that option by treating intermittent use as absence. A system built for variability loses resilience when every patch is assigned a permanent single purpose.

Permanent and mobile life were never clean opposites. Towns supplied grain, metalwork and markets; herders supplied animals, transport and products. Oasis farmers used caravan demand. Families combined settlement with seasonal movement. The desert economy worked by joining modes of life that later classifications separated.

The hydraulic city

A modern desert city can reach farther than a caravan or qanat. Dams store river flow gathered in distant mountains. Aqueducts and pipelines cross basins. Pumps lift groundwater. Treatment plants make wastewater usable again. Desalination converts seawater into municipal supply at the cost of energy, infrastructure and concentrated brine. Refrigeration and air conditioning shift heat management from architecture towards electricity.

The system works by separating local appearance from regional metabolism. A tree-lined street may be supported by snow falling hundreds of kilometres away. Food may arrive from wetter land. Electricity may arrive from a gas plant, solar field or interconnected grid. Waste leaves by pipe, lorry or atmosphere. The city is not outside the desert. It is a machine for importing missing flows and exporting some consequences.

Southern Nevada offers a clear local distinction. Water used indoors is treated and much of it returned to Lake Mead, allowing a return-flow credit against withdrawals. Water used on lawns and other outdoor landscaping is largely lost through evaporation and plant transpiration. Two households can draw the same amount at the meter while placing different net demands on the river. This is why conservation policy has targeted ornamental turf. The arrangement depends on treatment works, topography and Colorado River law, so it cannot be copied as a general desert-city rule.

Desalinated coastal cities solve a different shortage. They gain a supply that does not depend on local rain, but the plant depends on power, membranes, maintenance and a marine outlet for brine. Inland cities cannot copy the arrangement without dealing with transport and disposal. Technology changes where supply comes from and where the costs appear; it does not remove physical limits.

Stormwater adds a paradox. Streets and roofs prevent infiltration and accelerate runoff, so a city built against scarcity may also need large drains and flood basins. Water that could recharge ground becomes a hazard because hard surfaces move it too quickly. Capturing stormwater can help,

but contaminated runoff, limited storage and the rarity of events constrain the gain.

Heat follows the same built form. Dark surfaces and waste heat raise urban temperatures, increasing cooling demand and outdoor water loss. Shade trees can reduce heat while consuming water. Dense buildings reduce exposed area but may trap warm air. There is no design that maximises every objective. A desert city must choose where to spend water, land and energy, then make the trade visible.

The strongest systems use several supplies and several defences. They protect watersheds, price scarcity, repair leaks, recycle suitable wastewater, plan for drought and preserve flood routes. Diversity is valuable because each source fails differently. Dependence hidden beneath a single reliable tap is still dependence.

When renewal falls behind

A reserve can support growth long before depletion becomes visible. Wells deepen gradually. Pumps use more energy. Springs weaken. Wetlands contract. Farmers shift crops or abandon fields. By the time a regional decline is clear, buildings, debts and livelihoods may depend on continued withdrawal. The physical stock has become a political commitment.

Land degradation follows several routes. Repeated disturbance can break soil crust and expose fine sediment to wind. Grazing concentrated near fixed water can remove cover. Poor irrigation drainage can salt soil. Off-road vehicles and construction can alter runoff. Mining can change water chemistry or fragment habitat. Warming can raise evaporative demand and shift the threshold at which plants survive. These pressures interact, and the damaged state may persist after the original cause stops.

A satellite image can show more vegetation after rain without proving recovery. A restoration project can plant trees while lowering groundwater. A fence can protect one patch while blocking movement through a larger system. Good management follows the process that made the land productive: water pathways, soil stability, patch structure, mobility, rights

and time.

The decisive comparison is between renewable flow, stored reserve and reversibility. A storm can refill a pond. It may not refill a deep aquifer on a human timescale. A plant can return after one season. A broken pavement or lost endemic population may not. Scarcity becomes manageable when these differences are kept visible.

Feedbacks can push a system across a threshold. Loss of vegetation exposes soil. Erosion removes fine material and nutrients. Runoff becomes faster, giving new plants less chance to establish. Dust can darken snow elsewhere and alter melt. Falling groundwater weakens springs, concentrating wildlife and livestock around the few that remain. None of these loops operates everywhere, and some degraded land recovers when pressure is removed. The possibility of recovery does not make the path back cheap or quick.

Desertification policy has often failed by attacking the picture rather than the process. A line of trees may slow wind locally but die without water or displace grazing. Settling mobile people can make services easier to deliver while concentrating herds. Deep wells can reduce walking and then draw animals into one damaged radius. An intervention changes incentives and movement as well as soil. Its success must be judged across the wider system and over dry years, not from photographs taken after rain.

Recent aridification tightens these choices in many regions. Higher evaporative demand can turn the same rainfall into less usable moisture. The response is not to declare every dry place doomed. It is to update baselines, protect options and stop counting old reserves as new supply.

How we know

Deserts are difficult to measure because stations are sparse, events are local and long dry intervals tempt short studies. Rain gauges record what falls at one point. Stream gauges catch only monitored channels. Wells reveal water levels locally. Satellite gravity missions such as GRACE detect regional changes in stored water, while optical and radar sensors map vegetation, floods, dunes and surface disturbance. None describes the whole system alone.

Groundwater age and origin can be investigated through isotopes, dissolved gases and chemistry. Sediment, shorelines and salt deposits reconstruct former lakes and climates. Field experiments measure plant responses, soil crust, infiltration and animal physiology. Archaeology reveals routes, wells, fields and settlements that written records ignored.

Human knowledge is evidence too, though it must be used with the same care as instruments. Pastoralists, farmers and Indigenous communities may hold detailed records of water points, seasonal movement and ecological change. Their knowledge is situated rather than universal, as all observation is. The strongest picture comes from joining scales: a satellite trend, a field measurement, a historical record and the people living inside the system. The main uncertainty is seldom whether deserts are variable. It is whether the observation lasted long enough to witness the variability that matters.

What People Get Wrong

“A desert is hot and sandy”

Heat and sand make good cinema, so they became the definition.

Geography is less cooperative. Antarctica is a desert because it receives little precipitation. The Gobi has severe winters. The Namib can be cool and foggy near the coast. Sand seas are striking but occupy only part of most desert regions; rock, gravel, alluvial fans, dry lake beds and salt flats are often more extensive.

The correction matters because the wrong image hides the mechanism. Heat can intensify water loss, yet a desert is produced by a persistent moisture deficit. Sand is one sediment among many. Policies, surveys and expectations built around dunes miss cold deserts, stony rangelands, groundwater-fed wetlands and flood-prone channels. The desert is not a landform. It is a water condition expressed through many landforms.

The mistake also reverses risk. A cold desert can threaten through exposure and short growing seasons rather than extreme daytime heat. A rocky basin can flood faster than a dune field absorbs water. Knowing the surface and the source of aridity matters more than recognising a familiar postcard.

“Nothing lives there”

Desert life is sparse, hidden and badly timed for visitors. Much of it is underground, nocturnal, dormant, microscopic or concentrated around washes and springs. A survey taken in a dry year at midday can find little and still describe the ecosystem poorly. Seeds may wait in soil. Amphibians may remain buried. Biological crust can occupy the apparent gaps between plants.

The myth became persuasive because abundance is usually judged by continuous green cover. Deserts distribute productivity in patches and pulses. That does not make every desert equally diverse, and hyper-arid interiors can support little life. It means absence must be demonstrated across seasons, events and scales rather than inferred from a brown photograph. Calling land lifeless lowers the apparent cost of tracks, mines, roads and energy projects before anyone has measured what is there.

Some of the most consequential habitat is also the least extensive. A spring occupying a tiny fraction of a basin may support the only permanent water and species restricted to that outlet. A wash may carry the densest trees and provide a movement corridor. Area alone is a poor measure of ecological importance when resources are concentrated.

“Rain is always a blessing”

Water is valuable, but delivery matters. Intense rain on steep, sealed or sparsely vegetated ground can generate flash floods that carry sediment, destroy roads and kill people far from the storm. A downpour can scour plants, erode soil or trigger germination followed by lethal drying. In salt-affected ground, shallow water can raise salts towards roots as it evaporates.

The belief survives because annual totals frame rain as a missing commodity. Deserts need water, so more appears automatically better. The proper unit is a pathway: how fast it falls, where it runs, what infiltrates, what can be stored and who is exposed. A storm may damage a town, recharge an aquifer and feed a wetland in the same event. Blessing and hazard are not rival descriptions. They are different positions in the catchment.

This is why flood control can create a second problem. A channel straightened to move water away from buildings may reduce local infiltration and send a sharper peak downstream. A dam may protect one district while withholding sediment or recharge from another. The desirable outcome is not maximum capture or maximum drainage. It is a deliberate allocation of a rare, violent pulse.

“Camels store water in their humps”

The hump stores fat. It changes size as the animal uses and replaces that energy reserve. Metabolising fat produces some water, as metabolism does in other animals, but the hump is not a portable cistern. Camels survive dehydration through a suite of traits: reduced water loss, concentrated urine, dry faeces, tolerance of changing body temperature, recovery of moisture in the nose and rapid rehydration when water returns.

The tank story is memorable because it gives one visible structure one job. Biology is usually less tidy. The correction matters beyond trivia. Adaptation is integrated and costly. A camel still needs forage, water, rest and routes between them. Treating the animal as independent of supply turns a working human-animal network into a cartoon of limitless endurance.

It also hides the human achievement. Long-distance camel transport depended on breeding, loading, route knowledge, wells, fodder and organised stops. The animal widened the interval that could be crossed; it did not remove the interval. The caravan was a system of planning built around an adapted body.

“Nomads wander aimlessly”

Mobility in drylands is often highly organised. Herders move according to rain, forage, water, disease, markets, security, customary access and information from other people. Routes may include wet-season pasture, dry-season reserves and fallback areas used only in bad years. The pattern can change because the resource pattern changes.

Settled officials often read intermittent use as non-use. That made the myth administratively convenient: land that appeared empty could be fenced, cultivated, conserved or allocated. Movement can cause conflict and mobile systems can degrade land, so romanticising them is no improvement. The evidence rejects randomness, not criticism. The practical issue is whether mobility matches variability. Blocking a route can concentrate animals and worsen the degradation that settlement was meant to prevent.

A route can therefore be lost without a building being demolished. A border closes, a reserve excludes grazing, a farm takes a dry-season corridor or a borehole attracts permanent settlement. Each fixed change can remove an option that mattered only in exceptional years. Variability makes spare access look wasteful until the year it becomes essential.

“Desertification means the desert is marching forwards”

Advancing dunes can bury roads and fields, but they are one local process. Under the United Nations convention, desertification is the degradation of land function across drylands, driven by climatic variations and human activities. It can appear as lost vegetation, eroded soil, salinity, falling productivity, damaged water systems or ecological simplification without a grain of moving sand.

The marching-desert image became powerful because it offers a front line and a visible enemy. It also suggests that planting a barrier of trees will solve the problem everywhere. Real degradation is shaped by land rights, water, grazing, farming, infrastructure and climate at several scales. A damaged land system cannot be repaired by moving the colour green across a satellite image. Restoration has to recover function and fit the people who use the land.

Aridification is different again. It describes a long-term shift towards a drier climate. It can raise the risk of degradation without proving that degradation has occurred. Keeping climate, land condition and desert scenery separate prevents one map from being asked to answer three questions. It also makes successful recovery visible where management improves land despite a dry climate.

“Empty desert is harmless development land”

Low population and low vegetation cover are often converted into a claim of low value. The ground may instead contain slow-growing plants, cryptic soil crusts, migration routes, archaeological sites, flood pathways, endemic spring species, grazing rights or seasonal use. Damage can be hard to reverse because recovery is slow and water concentrates impacts.

This does not prohibit mines, roads, housing or solar energy. It changes the baseline. Development should be compared with the best alternative sites and with the full system it enters, including transmission, water demand and cumulative fragmentation. A disturbed industrial parcel and an intact desert plain may receive the same sunshine but carry different ecological costs. Empty is not an observation. It is a conclusion, and it is often reached before the survey begins.

Cumulative damage is easy to miss because each project occupies a small share of a large view. Roads split movement, power lines add access, wells draw from a common aquifer and vehicle tracks seed erosion. The question is not whether one footprint fills the desert. It is whether many footprints remove the connections on which a sparse system depends.

Use It

Follow the water, not the rainfall total

A rainfall number describes what crossed an imaginary horizontal surface. It does not tell you what became usable. Before judging scarcity, trace the route. Did the moisture arrive as gentle rain, a short cloudburst, snow, fog or river flow imported from another climate? Did it fall over a mountain catchment, a sealed city or loose sand? What evaporated, ran away, infiltrated, became polluted or remained beyond reach?

Then change the timescale. An annual mean can be built from a few rare storms separated by failure. A reservoir depends on sequences of years, not one average year. A plant may depend on rainfall reaching a particular soil depth during a narrow season. A settlement may depend on snow that falls far outside the desert. The useful questions are when the water arrived, where it travelled and how long its benefit remained. This prevents one impressive storm, one wet year or one green satellite image from being mistaken for a repaired water system.

Separate renewal from reserve

A working tap says nothing about the age of its source. Groundwater may be replaced by recent floods, slow mountain recharge, leakage from a river, or rain that fell under a wetter climate thousands of years ago. The same pumping rate can therefore be modest in one aquifer and depletion in another.

Ask for the level as well as the withdrawal. What is known about recharge, and at what scale? Are springs weakening while deep wells still produce? Does the estimate cover the whole basin or one monitored point? How quickly would the store recover if pumping stopped? Avoid the lazy division between renewable and fossil water as though every aquifer belongs cleanly to one class. Waters of different ages mix, and recharge can be small without being zero. The decision still turns on rates: withdrawal, replacement and the timescale on which users need the source to remain available.

Read pulses, thresholds and refuges

Desert systems rarely respond in proportion to an annual total. A light shower may wake microbes and evaporate before reaching roots. A larger event may fill a pool, recharge an alluvial channel or trigger breeding. Two storms close together can sustain growth that the same total scattered across months would not. A threshold separates no response from a large one.

Look for what must already be present when the event arrives. Seeds need the right temperature. Floodwater needs a route and somewhere to spread. A wetland needs enough spring discharge to maintain its chemistry. Animals need access to the temporary concentration of food or water. Also look for refuges that carry the system through the interval: deep soil moisture, shade, burrows, reserve pasture, seed banks and springs. Managing only the pulse can destroy the store or route that makes the pulse useful. Capturing every flood, for example, may protect one site while starving recharge or wetlands downstream.

Treat movement as infrastructure

In variable drylands, fixed resources are often individually insufficient. Wells, pasture, markets and shelter become workable when routes connect them. A pastoral corridor may leave no concrete trace, yet its closure can concentrate livestock around permanent water and remove a reserve used only during drought. A wildlife corridor may matter for a few weeks each year. A caravan route needed guides, credit, agreements and reliable staging points as much as it needed camels.

Map movement before allocating land. Who crosses after rain, during the dry season or when a usual source fails? Which rights make the route usable? Where do fences, borders, roads, mines or protected areas narrow the options? Mobility should not be romanticised. It can carry disease, intensify conflict or concentrate pressure in the wrong conditions. Its value is functional: it joins resources scattered across space and time. Removing that connection can turn low density from an adaptation into a trap.

Change scale before calling land empty

Sparse cover encourages hurried conclusions. A spring can occupy a tiny share of a basin and support species found nowhere else. A dry wash can be the main flood route, recharge zone and strip of woodland. The space between shrubs may be held by biocrust, roots and seed. A salt flat may connect to wetlands beyond the project boundary. Seasonal grazing or gathering may be invisible during a short survey.

Before treating a site as unused, observe it at more than one time and from more than one position. What happens after rain, at night, below ground, during migration or in a severe dry year? What passes through rather than staying? Which impacts appear only when several roads, wells or energy sites accumulate? Ask who supplied the baseline and whose use would leave the weakest physical mark. The answer does not grant every site immunity from development. It prevents zero value from being assumed because the value was dispersed, dormant or omitted from the map.

Restore the process, then test it through dry years

Green colour is an unreliable target. Trees can be planted where water cannot sustain them. A fence can thicken vegetation inside while shifting grazing pressure outside. Rain can make a damaged site look recovered for one season. Restoration should begin with the failed function.

Is runoff moving too quickly for infiltration? Has salt accumulated because drainage is blocked? Has a spring lost discharge? Is wind removing exposed soil? Are animals unable to reach seasonal pasture? The intervention should fit that mechanism: slowing water, protecting a crust, repairing drainage, changing access, reducing withdrawal or allowing recovery without planting. Measure the wider system, not the chosen plot alone. Include dry periods, because a project designed from wet-year photographs may vanish when it is needed. Success is a land system that retains soil, water, habitat or livelihood under its normal variability, not a greener image on the day of inspection.

The limits

A moisture-balance model explains a great deal, but it does not decide whose claim should prevail. Two users can understand an aquifer perfectly and still disagree over law, livelihood, history and fairness. A route can be ecologically useful and politically contested. A solar project can reduce emissions while damaging local habitat. Measurement clarifies the trade-off. It cannot choose the values.

Nor does one desert teach a universal method. Fog supports life in parts of the Namib and Atacama but says little about a continental interior. Southern Nevada's return-flow credits depend on treatment works, geography and Colorado River law. Results from a lithium basin or Mojave solar study cannot be copied to another basin without checking water pathways, species, rights and alternatives. Local knowledge has the same boundary. Long experience can reveal patterns that instruments miss, but it remains situated and can contain disagreement or unequal power.

Finally, adaptation has costs. CAM limits carbon gain. Dormancy sacrifices immediate growth. Mobility requires access and labour. Qanats require continual maintenance. Desalination requires energy and a place for brine. The fact that people or organisms survive does not prove the system is comfortable, just or indefinitely expandable.

The one thing to keep

Keep the renewal time.

When a desert looks empty, ask what is dormant, stored, moving or concentrated. Water may be underground. Life may be waiting for a storm. Pasture may matter in another season. A route may support the system while carrying nobody today. A salt flat may hold minerals, microbial life and connections to a wetland at once. The surface is a moment in a process whose important intervals may be separated by years.

Then ask how quickly each form of abundance returns. A pool may refill next season. A shrub may need decades after damage. A spring species

may have nowhere else to go. Deep groundwater may take centuries or millennia to replace. The same action can therefore be reversible at one scale and permanent at another.

That is what deserts should change in your view of the world. Scarcity does not mean nothing is present. It means value is unevenly distributed and often hidden from a short visit. Survival depends on matching use to timing, stores, refuges and routes. A green field may be an old aquifer passing through leaves. A reliable tap may be snow from another state or rain from another age. The decisive distinction is no longer empty against abundant. It is abundance that returns against abundance that can be spent once.

Terms

Aridity. A long-term shortage of available moisture relative to atmospheric demand. It describes climate, not a temporary emergency, and can result from low precipitation, high evaporative demand or both.

Aridity index. A ratio comparing precipitation with potential evapotranspiration. It gives a more useful measure of dryness than rainfall alone, though local water pathways and seasonality still matter.

Desert. A region with a severe persistent moisture deficit. Deserts include hot, cold, sandy, rocky, coastal and polar forms, so the word names a water condition rather than one appearance. Its boundaries are climatic gradients rather than walls.

Dryland. The wider family of water-limited regions, spanning climates from hyper-arid desert through arid and semi-arid zones to dry sub-humid land. Not every dryland is a desert, and much dryland supports grassland, farming and settlement. The distinction matters in land policy.

Drought. A period when water supply falls below the normal range or needs of a particular place or system. Drought is relative to a baseline; aridity is a long-term condition.

Hyper-arid. The driest climatic class, where precipitation is tiny relative to

evaporative demand. Hyper-arid cores may support little continuous vegetation while still containing episodic channels, microbes or groundwater-fed life.

Evapotranspiration. Water returned to the atmosphere through evaporation from surfaces and transpiration from plants. Potential evapotranspiration estimates atmospheric demand if water were available; actual loss is limited by supply.

Rain shadow. A dry region on the leeward side of mountains. Rising air rains on the windward slope, then descends, warms and becomes less humid after crossing the crest.

Subsidence. Downward movement of air. In subtropical high-pressure belts, descending air warms and suppresses cloud formation, helping create many of the world's major hot deserts.

Inversion. An atmospheric layer in which warmer air sits above cooler air, resisting vertical mixing. Coastal inversions can produce persistent fog while preventing the cloud growth needed for substantial rain.

Wadi. A channel or valley in an arid region that may remain dry for long periods and carry sudden floods after rain. Similar regional terms include arroyo and wash.

Alluvial fan. A fan-shaped deposit formed where a stream leaves a steep canyon, slows and drops sediment. Channels can shift across the fan, creating both recharge zones and flood hazards.

Playa. A flat-floored closed-basin depression that may hold a temporary lake. Repeated wetting, sedimentation and evaporation can produce clay pans, salt crusts and major dust sources.

Erg. A large sand sea containing dunes and sand sheets. Ergs dominate popular desert imagery but form only one part of the much wider range of desert surfaces.

Reg. A stony desert plain covered mainly by gravel or pebbles. The term is

used particularly in North Africa and contrasts with an erg's sand and a hamada's rock plateau.

Hamada. A high, hard, rocky desert surface or plateau from which much finer material has been removed. Hamadas show why the Sahara cannot be reduced to dunes.

Desert pavement. A closely packed surface layer of stones above finer sediment. It can protect soil from erosion and may take extremely long periods to re-form after heavy disturbance.

Biocrust. A living community of cyanobacteria, algae, fungi, lichens and mosses on or near the soil surface. Biocrusts can stabilise soil, alter water movement and contribute carbon or nitrogen. Their effects on infiltration vary by crust and soil.

CAM. Crassulacean acid metabolism, a photosynthetic pathway in which plants open stomata mainly at night, store carbon temporarily in acids and use it by day with pores closed.

Ephemeral. Lasting for a short time or active only after suitable conditions. Desert streams, pools, plants and animal activity may be ephemeral without being unimportant or accidental.

Pulse-reserve model. An ecological model in which irregular resource pulses, usually rain, trigger activity and organisms store gains in seed, tissue, fat, eggs or nutrients for the interval that follows. Responses differ by event size, depth, season and organism.

Aquifer. Permeable rock or sediment that stores and transmits groundwater. It is usually water held in pores or fractures, not an underground open lake.

Recharge. Water entering an aquifer. In deserts it may be concentrated beneath mountains, flood channels or rare storm paths, far from the wells that later withdraw it.

Fossil groundwater. Groundwater largely recharged under past climatic

conditions and renewed negligibly on management timescales. The label should be used cautiously because aquifers often contain waters of mixed ages.

Oasis. A productive place where water becomes accessible in a desert through a spring, shallow groundwater, flood diversion, wells or engineered supply. Its survival also depends on labour and rules.

Qanat. A gently sloping underground tunnel that intercepts groundwater at higher ground and carries it by gravity to settlement or fields. Vertical shafts provide construction and maintenance access.

Endorheic basin. A drainage basin with no surface outlet to the ocean. Water ends in lakes, wetlands, playas or groundwater and leaves mainly through evaporation or infiltration, concentrating salts. Some hold wetlands or mineral-rich brines.

Salinisation. Accumulation of soluble salts in soil or water. Irrigation and evaporation can raise salt levels around roots unless drainage removes them, reducing agricultural productivity.

Pastoralism. A livelihood organised around livestock and grazing land. In variable drylands, strategic mobility can connect scattered forage and water, though outcomes depend on access, stocking and institutions. Mobility is planned adaptation, not random wandering.

Desertification. Land degradation in arid, semi-arid and dry sub-humid areas resulting from climatic variations and human activities. It concerns lost function, not the simple advance of desert sand.

Go Deeper

For the quickest wider survey: Nick Middleton, *Deserts: A Very Short Introduction* (Oxford University Press, 2009). Middleton covers climate, landforms, wildlife, human use and environmental pressure in a compact volume written by a geographer who has worked across many of the world's deserts. Read it to widen the examples in this book without losing the central distinctions. Its brevity means individual regions pass quickly, but that is also its strength for a first step. Pay particular attention to the opening problem of definition: it prevents every later example from collapsing back into heat and dunes.

For a desert understood through its long-term residents: Gary Paul Nabhan, *The Desert Smells Like Rain: A Naturalist in O'odham Country*, 40th-anniversary edition (University of Arizona Press, 2022). Nabhan writes from long work with Tohono O'odham communities in the Sonoran Desert, joining water, crops, weather, language and observation. It is not a global account and should not be treated as one. Read it to see how a landscape dismissed as sparse becomes dense when knowledge is local, cumulative and lived. The anniversary edition adds a new preface reflecting on climate change, the border and the knowledge shared with him, while preserving the close observations that made the original influential.

For the physical system in depth: Julie J. Laity, *Deserts and Desert Environments* (Wiley-Blackwell, 2008). This is the serious next book on climate, hydrology, weathering, rivers, dunes, dust, soils and past environmental change. Laity treats deserts as linked systems rather than collections of dramatic landforms. It is a university-level text and denser than the other recommendations, but its diagrams and global range repay careful reading. Use it when you want the mechanism behind a particular claim made here. The chapters on water and landforms are especially useful for seeing why rare floods, groundwater and wind must be analysed together rather than assigned separate worlds.

For the argument about power and the idea of wasteland: Diana K.

Davis, *The Arid Lands: History, Power, Knowledge* (MIT Press, 2016).

Davis traces how European and colonial accounts often blamed mobile pastoralists and local land use for drying and degradation on evidence that did not justify the claim. Her critique is strongest as a warning about knowledge serving administration. It does not make human-caused degradation imaginary. Read it beside the physical science to understand why describing land as ruined, empty or mismanaged can be a political act before it becomes an environmental finding. Its subject is arid lands broadly, so use it as a challenge to categories and authority rather than as a field guide to every desert process.

Notes and Sources

The evidence for deserts is unusually sensitive to scale. A global aridity class, a basin water balance and a household water shortage can all be accurate while describing different things. The notes below therefore distinguish climatic desert, dryland, drought, aridification and desertification. Figures tied to particular observation periods or regions are labelled rather than presented as timeless global constants.

The Whole Thing in One Page and Why You Should Care

Definition and variety. The organising definition follows physical geography rather than the popular image of heat and dunes. Julie Laity's *Deserts and Desert Environments*, Nick Middleton's *Deserts* and Andrew Goudie's *Arid and Semi-Arid Geomorphology* treat deserts through moisture balance, landforms and regional variation. Antarctica is retained as a polar-desert example, but the narrative avoids one continental precipitation average because coastal and interior conditions differ greatly. The Sahara is described as the largest hot desert, preserving the distinction between the largest desert and the largest hot one.

Sand and surface cover. No universal percentage is used in the narrative because estimates depend on whether dunes, sand sheets and smaller

accumulations are combined, and on how desert boundaries are drawn. Remote-sensing and geomorphological literature nevertheless supports the correction that dunes occupy a minority of most desert regions. Rock, gravel, alluvium, clay and salt surfaces are extensive.

Lake Manly. The Death Valley opening uses National Park Service reports. The remnants of Hurricane Hilary delivered 2.2 inches of rain in August 2023. A further 1.5 inches arrived in February 2024. The temporary lake was reported at roughly six miles long, three miles wide and one foot deep, with kayaking possible for almost a month. These measurements describe the 2024 event, not a normal condition of Badwater Basin.

Drying lands. The recent trend discussion draws on *The Global Threat of Drying Lands*, a 2024 report of the Science-Policy Interface of the United Nations Convention to Combat Desertification. Comparing 1991 to 2020 with 1961 to 1990, it assigns an additional 4.3 million square kilometres to dryland classes and places their total above 40 per cent of land outside Antarctica. The manuscript uses the report's official area change and period labels rather than turning rounded class shares into false precision. This is an aridity classification based on long-term climate, not a direct measurement of degradation and not evidence of one uniform regional cause.

The Core Ideas

Aridity as a moisture balance. The distinction between precipitation alone and water balance follows standard use of the aridity index, which compares precipitation with potential evapotranspiration. Broad geographical classifications often group hyper-arid, arid, semi-arid and dry sub-humid climates as drylands. Article 1 of the UNCCD uses a narrower operational definition: a precipitation-to-potential-evapotranspiration ratio of 0.05 to 0.65, covering arid, semi-arid and dry sub-humid areas while excluding hyper-arid, polar and sub-polar regions. The difference is terminological rather than evidence that Antarctica is not climatologically a desert.

How deserts form. The account of subtropical descent, rain shadows, cold

currents, continental interiors and polar dryness synthesises standard atmospheric and physical geography in Laity, Middleton and Goudie. These mechanisms often overlap. Coastal fog deserts, for example, may combine cold currents, stable air and topography. The Green Sahara discussion is kept qualitative because the timing and intensity of African Humid Period changes vary across regions and archives.

Pulses and reserves. Imanuel Noy-Meir's 1973 review gave desert ecology its classic pulse-and-reserve model: irregular water inputs trigger production, and organisms store resources between events. Travis Huxman and colleagues and James Reynolds and colleagues refined the model by showing that pulse size, timing, soil depth and organism type alter the response. Ferran Garcia-Pichel and Osvaldo Sala later extended the logic to microorganisms. The book uses the model as a guide, not a claim that every desert response is identical.

Floods, infiltration and groundwater. Philip Stoffer's United States Geological Survey report on Mojave desert landforms, together with the hydrology and geomorphology syntheses by Laity and Thomas, supports the account of rare storms producing destructive floods while also contributing recharge in suitable settings. Recharge is highly uneven. Some floodwater evaporates or exits a basin; some enters shallow alluvium; some reaches deeper aquifers only under particular geology and event size. The term fossil groundwater is used for water with negligible modern replacement on human management timescales, not as a label for all deep groundwater.

Plant water economy. Karolina Heyduk's 2022 review in *Plant Physiology* supports the CAM explanation. CAM plants take up much of their carbon dioxide at night, store it as malic acid in vacuoles and release it internally during the day, reducing water loss from daytime gas exchange. CAM varies among species and conditions, so the narrative avoids implying that every succulent or desert plant follows the same schedule.

Animal physiology. Knut Schmidt-Nielsen's *Desert Animals* remains a foundational comparative treatment of heat and water balance and supports

the camel mechanisms used here: fat stored in the hump, tolerance of changing body temperature, concentrated urine, reduced respiratory water loss and rapid rehydration. The common statement that a camel carries a tank of water in its hump is false. Exact endurance figures are omitted because they change with heat, workload, diet, health and species.

Desert surfaces and living crust. The sections on alluvial fans, dunes, desert pavement and dust use Stoffer's USGS report and the geomorphology texts by Laity, Goudie and David Thomas. Biological soil crust is described from the synthesis edited by Jayne Belnap and Otto Lange and from dryland microbial ecology. Crust composition and function vary, and the book therefore avoids one recovery time or one universal effect. Vehicle scars can persist for decades or longer in some settings, while other surfaces recover faster.

Saharan dust. Hongbin Yu and colleagues used CALIPSO observations from 2007 to 2013 to estimate that African dust supplied about 0.022 teragrams, or 22,000 tonnes, of phosphorus to the Amazon basin per year, with a stated uncertainty range of 6,000 to 37,000 tonnes. The study found this input comparable to estimated hydrological phosphorus losses. The book states the observation period and uncertainty because transport varies from year to year and later studies use different models, periods and basin definitions.

Human adaptation and pastoral mobility. Gary Paul Nabhan's account of O'odham country supports the cautious Sonoran example of scarce-water use, cultivated crops, edible wild plants and place-based knowledge. It is retained as one lived desert relationship, not a global template. Matthew Turner and Eva Schlecht's 2019 critical review is used to resist two opposite simplifications: that pastoral movement is random, and that all movement is automatically sustainable. Their review finds movement shaped by livestock nutrition, water, security, markets, labour and access, with wide variation between systems.

Household water labour. The 2023 WHO and UNICEF Joint Monitoring Programme report uses individual-level data for households where water is

collected away from the premises. Across countries with relevant data, women and girls were responsible for collection in seven out of ten such households. The manuscript retains the unequal distribution of labour without presenting it as a desert-only pattern or a rule for every household.

Caravans and trade. UNESCO's World Heritage record for Timbuktu and Volume IV of the *General History of Africa* support the account of desert routes linking West Africa, North Africa and Mediterranean markets. The list of traded goods includes salt, gold, cloth, books and enslaved people. The caravan is presented as an organised system of water points, guides, credit, protection and staging rather than as a camel alone defeating geography.

Qanats and Yazd. UNESCO's World Heritage records for the Persian Qanat and the Historic City of Yazd support the description of gently sloping underground channels, vertical access shafts, gravity flow, communal maintenance, windcatchers, courtyards and thick earthen walls. Qanats occur under different names across a wide region and have diverse dates and designs. The Persian example is retained as a documented system, not claimed as the only origin of underground water galleries.

Southern Nevada. The return-flow example comes from the Southern Nevada Water Authority. Treated Colorado River water returned through the Las Vegas Wash to Lake Mead earns return-flow credits, allowing a corresponding further withdrawal within the legal accounting system. Indoor use is much more recoverable than outdoor irrigation, where water is largely consumed by evapotranspiration. The example is setting-specific because it depends on geography, treatment infrastructure and interstate river law.

Recent groundwater trends. Scott Jasechko and colleagues analysed in situ groundwater-level trends from about 170,000 monitoring wells in 1,693 aquifer systems in more than 40 countries, with records spanning at least eight years. The covered countries account for about 75 per cent of global groundwater withdrawals, but the wells are not a random global sample and coverage is uneven. The study distinguishes groundwater level from total recoverable storage, shows strong local variability and reports both widespread rapid decline in cultivated dry regions and documented cases of

slowed or reversed decline after intervention. GRACE remains useful for broad storage trends but is too coarse for local management. The earlier 2015 GRACE studies are retained to show regional trend detection and uncertainty in total storage, not to estimate one local aquifer.

Lithium brines. Alexander Kirshen and colleagues' 2025 study calculated modern freshwater inflows for 28 active or prospective lithium-producing closed basins in Chile, Argentina and Bolivia. The authors found inflows of 2 to 33 millimetres per year and substantial overestimation by two commonly used global hydrological models. Direct streamflow measurements were available for only three basins. Daniel Corkran and colleagues' 2025 study then used groundwater-flow models representing three closed-basin endmembers, together with observations from two mine settings, to examine how abstraction location and fluid density affect wetland discharge. The manuscript uses these studies to distinguish freshwater from brine pathways and to require basin-specific assessment. It does not repeat their modelled effect percentages as universal impact numbers or treat either study as proof for every salar. Both papers acknowledge funding or support from BMW Group and BASF; their findings are used with explicit method and transportability limits.

Solar development. Jason Kreidler and colleagues' Mojave case study shows how conservation planning can compare renewable-energy development with biodiversity values across a region. The book uses the case to establish that utility-scale solar has land, road, drainage and fragmentation effects and that siting changes the trade-off. It does not imply that Mojave species distributions or mitigation options apply unchanged elsewhere.

Desertification. Article 1 of the UNCCD covers degradation within arid, semi-arid and dry sub-humid areas and attributes it to climatic variations and human activities. James Reynolds and colleagues' 2007 synthesis and Diana Davis's environmental history support the rejection of a single advancing-desert model and of one universal cause. The book preserves room for documented damage from grazing, cultivation, irrigation, fire,

extraction and climate while resisting automatic blame assigned from a distant view.

Operating sequence and evidence

The operating sequence follows water from atmospheric supply through runoff, infiltration, storage, biological response and human allocation. Laity and the edited volume *Arid Zone Geomorphology* support the physical sequence. Noy-Meir, Huxman and Reynolds support the ecological response. The human sections combine documented technologies and practices rather than presenting one ideal desert society.

The discussion of endorheic basins and salinity follows basic mass balance: water can leave a closed basin chiefly through evaporation or subsurface leakage while dissolved material remains. Irrigation salinity depends on water quality, evaporation, soil, crop, drainage and management, so green production is neither evidence of permanent success nor proof of inevitable failure.

How we know. Rain gauges, stream gauges, satellite observations, isotope hydrology, sediment records, field experiments, archaeology and local knowledge operate at different scales and have different blind spots. The final paragraph's argument for joining them is methodological synthesis. No source is treated as universal merely because it is quantitative, and local knowledge is not treated as universal merely because it is long-held.

What People Get Wrong and Use It

The seven corrections draw on the same evidence above. The camel correction is anatomical and physiological. The nomad correction is institutional and ecological. The desertification correction is definitional and causal. The development correction is a baseline problem: apparently sparse land may contain routes, seasonal habitat, groundwater dependence or living soil that a snapshot misses.

The six lenses in Use It remain inside the subject. They ask readers to trace water pathways, distinguish renewal from reserve, identify pulses and

refuges, map mobility, change observation scale before declaring land empty, and restore function rather than colour. Their limit is deliberate: a moisture-balance model explains much, but it does not replace history, law, culture, power or place-specific evidence.

Terms and Go Deeper

Terms follow common usage in physical geography, ecology, hydrology and dryland studies. Where a term has a formal treaty meaning, such as desertification, that meaning is stated. The four Go Deeper editions and publication details were checked against publisher or library records on 2 September 2026.

Bibliography

Institutional and reference sources

British Antarctic Survey. “Antarctic Factsheet and Geographical Statistics.”

British Antarctic Survey. Accessed 2 September 2026.

National Park Service. “Rare Opportunity to Kayak in Death Valley National Park.” 16 February 2024.

Southern Nevada Water Authority. “Where Your Water Comes From.”

Accessed 2 September 2026.

United Nations Convention to Combat Desertification. *United Nations Convention to Combat Desertification in Those Countries Experiencing Serious Drought and/or Desertification, Particularly in Africa*. Paris, 1994.

Vicente-Serrano, Sergio M., Narcisa G. Pricope, Andrea Toreti, Enrique Morán-Tejeda, Jonathan Spinoni, Anahi Ocampo-Melgar, Emma Archer, Arona Diedhiou, Tayebah Mesbahzadeh, Nijavalli H. Ravindranath, Roger S. Pulwarty, and Sara Alibakhshi. *The Global Threat of Drying Lands: Regional and Global Aridity Trends and Future Projections*. A Report of the Science-Policy Interface. UNCCD-SPI Technical Series No. 9. Bonn: United Nations Convention to Combat Desertification, 2024.

Stoffer, Philip W. *Desert Landforms and Surface Processes in the Mojave National Preserve and Vicinity*. USGS Open-File Report 2004-1007. Reston, VA: United States Geological Survey, 2004. doi: 10.3133/ofr20041007.

UNESCO World Heritage Centre. "Historic City of Yazd." World Heritage List, no. 1544.

UNESCO World Heritage Centre. "Timbuktu." World Heritage List, no. 119.

UNESCO World Heritage Centre. "The Persian Qanat." World Heritage List, no. 1506.

United Nations Children's Fund and World Health Organization. *Progress on Household Drinking Water, Sanitation and Hygiene 2000-2022: Special Focus on Gender*. New York: UNICEF and WHO, 2023.

Books

Belnap, Jayne, and Otto L. Lange, eds. *Biological Soil Crusts: Structure, Function, and Management*. Ecological Studies 150. Berlin and Heidelberg: Springer, 2003.

Davis, Diana K. *The Arid Lands: History, Power, Knowledge*. Cambridge, MA: MIT Press, 2016.

Goudie, Andrew S. *Arid and Semi-Arid Geomorphology*. Cambridge: Cambridge University Press, 2013.

Laity, Julie J. *Deserts and Desert Environments*. Chichester: Wiley-Blackwell, 2008.

Niane, D. T., ed. *General History of Africa, Volume IV: Africa from the Twelfth to the Sixteenth Century*. Paris: UNESCO, 1984.

Middleton, Nick. *Deserts: A Very Short Introduction*. Oxford: Oxford University Press, 2009.

Nabhan, Gary Paul. *The Desert Smells Like Rain: A Naturalist in O'odham Country*. 40th Anniversary ed. Tucson: University of Arizona Press, 2022.

Schmidt-Nielsen, Knut. *Desert Animals: Physiological Problems of Heat and Water*. Oxford: Clarendon Press, 1964.

Thomas, David S. G., ed. *Arid Zone Geomorphology: Process, Form and Change in Drylands*. 3rd ed. Chichester: Wiley-Blackwell, 2011.

Research articles

Corkran, Daniel B., David F. Boutt, Lee Ann Munk, Brendan J. Moran, Sarah V. McKnight, Jordan Jenckes, and Alexander Kirshen. "Density Constrains Environmental Impacts of Fluid Abstraction in Closed-Basin Lithium Brines." *Water Resources Research* 61, no. 9 (2025): e2024WR039511. doi: 10.1029/2024WR039511.

Garcia-Pichel, Ferran, and Osvaldo Sala. "Expanding the Pulse-Reserve Paradigm to Microorganisms on the Basis of Differential Reserve Management Strategies." *BioScience* 72, no. 7 (2022): 638-650. doi: 10.1093/biosci/biac036.

Heyduk, Karolina. "Evolution of Crassulacean Acid Metabolism in Response to the Environment: Past, Present, and Future." *Plant Physiology* 190, no. 1 (2022): 19-30. doi: 10.1093/plphys/kiac303.

Huxman, Travis E., Keirith A. Snyder, David T. Tissue, A. Joshua Leffler, Kiona Ogle, William T. Pockman, Darren R. Sandquist, Daniel L. Potts, and Susan Schwinning. "Precipitation Pulses and Carbon Fluxes in Semiarid and Arid Ecosystems." *Oecologia* 141, no. 2 (2004): 254-268. doi: 10.1007/s00442-004-1682-4.

Kirshen, Alexander B., Brendan J. Moran, Lee Ann Munk, Aeon A. Russo, Sarah V. McKnight, Jordan Jenckes, Daniel B. Corkran, Magdalen Bresee, and David F. Boutt. "Freshwater Inflows to Closed Basins of the Andean Plateau in Chile, Argentina, and Bolivia." *Communications Earth & Environment* 6 (2025): 177. doi: 10.1038/s43247-025-02130-6.

Jasechko, Scott, Hansjörg Seybold, Debra Perrone, Ying Fan, Mohammad Shamsudduha, Richard G. Taylor, Othman Fallatah, and James W. Kirchner. "Rapid Groundwater Decline and Some Cases of Recovery in

Aquifers Globally.” *Nature* 625 (2024): 715-721. doi: 10.1038/s41586-023-06879-8.

Kreidler, Jason, Carrie A. Schloss, Oliver Soong, Lee Hannah, and Frank W. Davis. “Conservation Planning for Offsetting the Impacts of Development: A Case Study of Biodiversity and Renewable Energy in the Mojave Desert.” *PLOS ONE* 10, no. 11 (2015): e0140226. doi: 10.1371/journal.pone.0140226.

Noy-Meir, Imanuel. “Desert Ecosystems: Environment and Producers.” *Annual Review of Ecology and Systematics* 4 (1973): 25-51. doi: 10.1146/annurev.es.04.110173.000325.

Reynolds, James F., Paul R. Kemp, Kiona Ogle, and Roberto J. Fernández. “Modifying the ‘Pulse-Reserve’ Paradigm for Deserts of North America: Precipitation Pulses, Soil Water, and Plant Responses.” *Oecologia* 141, no. 2 (2004): 194-210. doi: 10.1007/s00442-004-1524-4.

Reynolds, James F., D. Mark Stafford Smith, Eric F. Lambin, B. L. Turner II, Michael Mortimore, Simon P. J. Batterbury, Thomas E. Downing, Hadi Dowlatabadi, Roberto J. Fernández, Jeffrey E. Herrick, Elisabeth Huber-Sannwald, Hong Jiang, Rik Leemans, Tim Lynam, Fernando T. Maestre, Miguel Ayarza, and Brian Walker. “Global Desertification: Building a Science for Dryland Development.” *Science* 316, no. 5826 (2007): 847-851. doi: 10.1126/science.1131634.

Richey, Alexandra S., Brian F. Thomas, Min-Hui Lo, James S. Famiglietti, Sean Swenson, and Matthew Rodell. “Uncertainty in Global Groundwater Storage Estimates in a Total Groundwater Stress Framework.” *Water Resources Research* 51, no. 7 (2015): 5198-5216. doi: 10.1002/2015WR017351.

Richey, Alexandra S., Brian F. Thomas, Min-Hui Lo, John T. Reager, James S. Famiglietti, Katalyn Voss, Sean Swenson, and Matthew Rodell. “Quantifying Renewable Groundwater Stress with GRACE.” *Water Resources Research* 51, no. 7 (2015): 5217-5238. doi: 10.1002/2015WR017349.

Turner, Matthew D., and Eva Schlecht. "Livestock Mobility in Sub-Saharan Africa: A Critical Review." *Pastoralism* 9 (2019): 13. doi: 10.1186/s13570-019-0150-z.

Yu, Hongbin, Mian Chin, Tianle Yuan, Huisheng Bian, Lorraine A. Remer, Joseph M. Prospero, Ali H. Omar, David Winker, Yuekui Yang, Yan Zhang, Zhibo Zhang, and Chun Zhao. "The Fertilizing Role of African Dust in the Amazon Rainforest: A First Multiyear Assessment Based on CALIPSO Lidar Observations." *Geophysical Research Letters* 42 (2015): 1984-1991. doi: 10.1002/2015GL063040.