

THE IN A HURRY SERIES

Sleep *in a Hurry*

Rhythms, routines, and real rest

Max Brocklesby

BOOKSINAHURRY.COM

Contents

The Whole Thing in One Page 4

Why You Should Care 5

The Core Ideas 7

How It Actually Works 21

What People Get Wrong 31

Use It 36

Terms 40

Go Deeper 44

Notes and Sources 46

Bibliography 51

The Whole Thing in One Page

Sleep looks like absence. The eyes close, movement dwindles, the outside world loses access, and several hours disappear. That appearance supports two bad ideas: that sleep is blank time, and that a disciplined bedtime should be able to summon it on command.

Sleep is better understood as a timed, gated biological state. Two established processes decide when it is likely. Homeostatic pressure rises with time awake and falls during sleep. Circadian timing marks the body's internal day and night. They can disagree, which is why exhaustion and alertness can coexist during jet lag, shift work or an evening second wind.

For a practical mental model, two further conditions matter. Arousal must fall far enough for the transition to occur, and the body must be able to maintain sleep once it begins. Threat, pain, noise, learned vigilance and an overlong effort to force sleep can hold the first gate open. Airway obstruction, leg sensations, temperature, caring duties and unstable sleep-wake control can keep reopening the second. The two-process model explains timing. It does not explain every broken night.

Once sleep begins, the night changes state repeatedly. Non-REM and REM sleep recur in cycles, with deep slow-wave sleep weighted towards the early night and REM becoming more prominent later. Brain activity, muscle tone, breathing, heart rate and temperature control change with the stage. Equal durations can therefore contain different nights.

Healthy sleep is not perfect isolation. The sleeper must disconnect enough for stable cycles while retaining protective routes back to wakefulness. An arousal can restore breathing or answer a child's cry; hundreds of unnecessary arousals can dismantle continuity. The difference lies in cause, frequency and consequence.

Sleep supports waking performance, learning, memory, emotional regulation, immune function, metabolism and tissue maintenance. No single function explains all of it. The popular claim that sleep washes toxins from

the brain is especially unsettled. Animal studies using different methods disagree about clearance, and a 2026 human crossover study offered biomarker evidence without settling the mechanism because its inference relied on an investigational device and model. This is an active research field, not a consumer detox promise.

Real rest is wider than duration. Timing, regularity, continuity, satisfaction, daytime alertness, freedom from disruptive disorders and the night's internal structure all matter. Seven or eight hours can be mistimed, fragmented or repeatedly obstructed. A shorter night can be adequate for an unusual person and inadequate for most others. The result appears in daylight, but self-assessment also becomes less reliable as restriction accumulates.

Routines work indirectly. Light, rise time, activity, naps, caffeine, alcohol, temperature, opportunity and what the bed has come to predict change the conditions around sleep. They should reduce conflict, not become an evening examination. They also have limits. No bedtime routine can open a collapsing airway, remove a night shift, reliably stop menopausal hot flushes, silence a neighbouring nightclub or replace cognitive behavioural therapy for chronic insomnia.

The right question after a poor night is therefore not, "Why did I fail?" It is: which condition was missing, what pattern would distinguish it, and does the answer require a habit, a household change, a different schedule or clinical care?

That is the book.

Why You Should Care

In a controlled laboratory study, forty-eight healthy adults were assigned either eight, six or four hours in bed for fourteen consecutive nights, while another group underwent total sleep deprivation. The four-hour and six-hour groups accumulated dose-dependent lapses and cognitive deficits. Their subjective sleepiness rose early, then changed far less than their measured performance. They were losing the ability to judge the loss.

The experiment was small, artificial and limited to adults aged twenty-one to thirty-eight. It does not predict the exact impairment of every parent, clinician or night worker. It establishes a more important problem: the brain that needs judging is also the judge. Feeling accustomed to short sleep cannot certify that attention, reaction and decision-making have adapted with it.

Daytime sleepiness also differs from fatigue. Sleepiness pulls towards dozing; fatigue can feel like depletion without an ability to sleep. Pain, anaemia, infection, medication and low mood can produce fatigue, so a full night or reassuring tracker score cannot explain every exhausted day.

Sleep therefore matters before it becomes a medical complaint. It shapes driving, work, learning, training, appetite, mood and relationships. A person can become slower, more irritable or more error-prone while attributing each failure to workload, character or bad luck. The lost hour appears to have been converted into productive time because its cost is paid later, in smaller pieces, by the person making the calculation.

Sleep also exposes the limits of self-control. Modern schedules move meals, extend light, rotate shifts, cross time zones and negotiate waking with alarms. Internal clocks continue to use light and molecular cycles. When social time and biological time diverge, the result can be sleepiness during required wakefulness and alertness during the available night. A late chronotype may be harmless on one timetable and punishing on another.

The safety consequences are not private. A driver cannot bargain with a microsleep. Clinicians, pilots, warehouse workers and machine operators cannot replace sleep with professionalism near the circadian low. Employers, schools and transport systems shape sleep through shift design, start times, workload and whether fatigue can be reported without penalty. What looks like poor personal discipline may have been scheduled by somebody else.

Advice also changes direction according to the problem. More time in bed helps when opportunity is too short, yet can maintain some forms of insomnia by spreading sleep across a longer period of wakeful effort.

Alcohol can shorten the wait for unconsciousness while reducing REM sleep and disturbing later sleep. A night mode can reduce part of a screen's light signal while leaving another hour of work or novelty untouched. Catch-up sleep can help without making repeated weekday restriction free.

Then comes the market. Watches, rings, mattresses, lamps, supplements and apps promise control over a state that cannot be commanded directly. Some reveal useful patterns. None can turn a proprietary stage estimate into a clinical diagnosis. A score can make a good sleeper anxious and give a poor sleeper false reassurance. Measurement earns authority only when it matches the question.

This book replaces ritual with a model. You will understand how pressure and timing set the window, why arousal and bodily integrity decide whether the window holds, what changes across a cycle, what sleep is known to support, why healthy sleep has several dimensions, how routines alter the system, how chronic insomnia becomes self-maintaining, and when the problem has left the self-help category.

The limits stay visible. Caring duties, pain, hot flushes, disability, noisy housing, financial pressure and night work can remove opportunities that no ideal routine restores. Sleep is robust enough to survive an imperfect evening and vulnerable enough to punish a repeated mismatch. The task is to use the control you have, identify the control you lack and recognise when the body's refusal is evidence.

The Core Ideas

Sleep Is Permitted, Not Commanded

The common model gives sleep one cause: fatigue. Stay awake long enough, become tired enough, then fall asleep. Fatigue matters, but it cannot explain why someone who fought to remain awake at four in the afternoon can become alert at ten, or why jet lag can produce exhaustion and insomnia at the same time.

The first part of a better model is homeostatic sleep pressure. The need for sleep tends to rise across wakefulness and fall during sleep. Adenosine signalling participates in that regulation, but there is no single chemical fuel gauge that fills evenly from morning to night. Caffeine blocks adenosine receptors and makes pressure less perceptible. It changes the message, not the amount of wakefulness already incurred.

The second part is circadian timing. Cells throughout the body contain molecular clocks, coordinated by the suprachiasmatic nucleus in the brain. Specialised retinal cells report environmental light, helping align internal time with the day. The clock then coordinates rhythms in alertness, temperature, hormone release and many other processes. Melatonin is one signal of biological night. It can help shift timing in selected circumstances, yet it is neither a sleeping pill made by the body nor the switch that turns consciousness off.

Sleep becomes likely when pressure is high and the circadian system permits it. The forces can oppose each other. A nap lowers pressure. An evening wake-maintenance signal can temporarily conceal tiredness. An all-nighter creates severe pressure while the clock still produces periods of relative alertness. A traveller can arrive physically spent while the internal afternoon continues. A night worker can sleep after a shift, then wake early because daylight and circadian wake promotion are pushing in the opposite direction.

Those two processes are the strongest compact model of normal timing. They are not the whole event. Arousal determines whether the transition can occur. Threat, excitement, pain, work, clock-checking and learned expectation can maintain the waking state even when pressure and timing favour sleep. Opportunity determines whether the window exists at all. A parent caring for an infant and a worker finishing at 2 a.m. may have sound physiology and inadequate access.

A final condition is continuity. Once sleep begins, breathing, movement, temperature, neurological stability and the environment determine whether it remains consolidated. Obstructive sleep apnoea can provoke repeated

protective arousals. Restless legs can prevent stillness. Menopausal hot flushes, pain, medication, noise and caring duties can keep reopening the night. These failures may feel identical at breakfast and require different responses.

This broader model explains why bedtime is a poor place to begin many repairs. Much of the night has already been prepared. Light has nudged the clock. Wakefulness and naps have altered pressure. Caffeine has changed perception. Work has carried arousal towards bed. The schedule has either protected opportunity or spent it. A bedroom can remove noise and glare, but it cannot reverse every decision and constraint of the preceding day.

Direct control remains weak. Nobody can choose the minute at which N2 begins. Indirect control can be substantial. You can shape pressure, timing, opportunity, arousal and access to treatment. The useful questions are therefore wider than “How tired am I?” What time does the body think it is? Is there enough opportunity? What is holding wakefulness in place? What is interrupting sleep after it starts? The answer decides whether the next move is a routine, a schedule change, treatment or no intervention at all.

A Night Is a Sequence, Not a Block

In 1953, Eugene Aserinsky and Nathaniel Kleitman reported periods during sleep in which the eyes moved rapidly and brain activity changed. The discovery did not find a small decorative feature. It broke the idea that sleep was one uniform descent into inactivity.

Clinical sleep studies now divide sleep into rapid eye movement sleep, REM, and three non-REM stages. N1 is the unstable border between wakefulness and sleep. N2 is established light sleep and usually occupies the largest share of an adult night. Its electrical signatures include sleep spindles, brief bursts of rhythmic activity, and K-complexes, large waves that may help the sleeping brain respond selectively to the environment without waking fully. N3 is slow-wave sleep, the deepest conventional stage. REM combines active brain patterns and vivid dreaming with marked reduction of skeletal muscle tone.

These states recur in cycles, often four to six times in a night. The first cycle is not a template repeated with mechanical precision. Deep N3 sleep is concentrated towards the early part of the night, especially when sleep pressure is high. REM periods usually lengthen later. Brief awakenings become more common with age and can occur in healthy sleep. The architecture changes across infancy, adolescence, adulthood and old age, and it changes after sleep loss, illness, alcohol, medication and circadian displacement.

Stage labels also compress continuous biology into useful boxes. A scorer assigns a thirty-second epoch to the pattern that dominates, even though the brain may be crossing a boundary within it. Two laboratories can disagree at the margins without either seeing a different night. This is not a flaw that makes staging worthless. It is a reminder that the labels are maps of changing electrical and muscular patterns, not sealed chambers through which every sleeper passes on a perfect schedule.

This matters because “seven hours” does not identify what happened inside them. Two nights with equal duration can differ in continuity, timing, breathing, movement and stage distribution. A person with obstructive sleep apnoea may spend enough time in bed while repeated airway obstruction fragments the night. Alcohol may bring earlier unconsciousness and then more disruption. A new parent may accumulate several hours in pieces and feel nothing like someone who slept the same total continuously.

Trimming a night from one end is not neutral, but the effect is subtler than a first-half and second-half rule. Rising early reliably removes some of the REM-weighted later opportunity. Delaying bedtime while keeping the alarm shortens the night too, yet high homeostatic pressure can prioritise slow-wave activity. In restriction studies, REM and N2 may fall more while slow-wave activity is partly preserved or intensified. Circadian timing and accumulated loss still alter the pattern. The safe conclusion is not that early sleep is “for the body” and late sleep “for the brain”. Both NREM and REM support several functions, and neither is guaranteed by the clock alone. A full night protects a sequence whose proportions adapt to recent sleep

history.

Consumer discussions often turn that sequence into a league table. Deep sleep becomes premium, REM becomes creative, and light sleep looks like wasted space. The hierarchy is false. N2 is not filler. Spindles are implicated in learning and sensory protection. REM is neither the only dreaming stage nor a nightly certificate of emotional health. Deep sleep is important but cannot be ordered by staring at a score. Healthy sleep contains all these states because the sequence, not one winning stage, is the product.

A useful picture is a railway line rather than a tank being filled. Duration tells you how long the service ran. Architecture tells you which stations it reached, in what order, and how often it was stopped. You need the timetable as well as the total.

Rest Is Active, but No Single Job Explains It

Sleep-like states occur across diverse animal lineages despite an obvious cost. A sleeping animal is less able to find food, mate, defend territory or detect danger. Forms vary with ecology, and defining sleep outside familiar mammals is not always straightforward. Their repeated retention is consistent with benefits that outweigh vulnerability, but no single function explains every form of sleep.

The first function is visible the next day. Repeated restriction impairs sustained attention, reaction time and working accuracy. The dangerous feature is that subjective adaptation can outrun objective performance. A person may become familiar with sleepiness while lapses continue to accumulate. Confidence is therefore a poor safety test after a run of short nights.

Learning supplies another function. Wakefulness acquires information; sleep helps stabilise, reorganise and integrate parts of it. Different tasks recruit different features, so “sleep consolidates memory” is a heading rather than a complete mechanism. Slow-wave activity, spindles, REM, the timing of learning and the timing of recall can each matter. Sleep does not file every experience in a permanent cabinet. It changes which traces endure,

strengthen or connect.

Emotion sits inside the same two-way system. Poor sleep can increase negative affect and weaken regulation, while anxiety, depression and stress can damage sleep. One short night does not create a psychiatric disorder, and treatment of sleep is no universal cure. The stronger claim is that sleep participates in the processes by which the brain assigns salience, controls reaction and recovers from waking demand.

The sleeping body changes its allocation of work. Autonomic activity, hormone release, glucose handling, immune signalling and tissue processes follow organised patterns. Experimental restriction can alter insulin sensitivity, appetite, inflammation and immune responses, though the size and duration depend on dose, population and protocol. Long-term associations between short sleep and disease are important but frequently observational. Illness, work, stress, medicines and social conditions can affect both sleep and health. Chronic shortage deserves attention without turning an extra hour into guaranteed protection from a named disease.

Brain clearance shows how quickly one mechanism can become the whole story. A 2013 mouse experiment reported greater interstitial exchange and metabolite clearance during sleep. In 2024, another group measuring tracer departure in male mice reported reduced clearance during sleep and anaesthesia. A 2025 mouse study linked NREM noradrenaline, vascular and cerebrospinal-fluid oscillations to glymphatic flow. In 2026, a randomised crossover study of thirty-nine people found morning plasma amyloid beta and tau patterns consistent with sleep-active clearance, but the inference depended on an investigational device and a compartment model. These studies differ in species, tracer, route and outcome. They support active research into fluid transport, not a settled nightly detox mechanism or consumer optimisation ritual.

Sedation therefore cannot be judged by speed of unconsciousness. A drug may reduce responsiveness while changing architecture, continuity or breathing. Alcohol is the familiar example. At higher doses it can shorten sleep latency. Even lower doses can delay REM onset and reduce REM

duration, while effects on total sleep time, efficiency and later wakefulness vary across studies. Unconsciousness describes loss of report. Rest describes the quality and consequences of an organised state.

No single function carries the entire explanation because sleep coordinates many forms of maintenance across several scales. That is why one “best stage”, supplement or tracker metric cannot stand in for the night. Sleep is part of the production of competent wakefulness. Cut it, and the bill arrives during the hours that seemed to have been gained.

Healthy Sleep Is More Than a Number

“Eight hours” survives because it is memorable, schedulable and close to a useful centre for many adults. It becomes misleading when treated as a biological pass mark or the complete definition of rest.

The joint consensus of the American Academy of Sleep Medicine and Sleep Research Society recommends that adults regularly obtain seven or more hours to support health. This is a population boundary, not an individual ideal and not permission to assume that seven is enough for every person. Some adults need more. A rare inherited short sleeper may function well on unusually little. Most people sleeping five hours are not displaying an exotic genetic advantage. They are sleeping five hours.

Need changes across life and circumstance. Infants distribute sleep across day and night. Adolescents often shift later while retaining a substantial sleep requirement, which makes an early school timetable biologically expensive. Pregnancy, illness, recovery and heavy training can alter sleep need, opportunity or continuity without yielding one universal extra-hour rule. Menopause may bring night waking through vasomotor symptoms, mood change and other interacting causes. Older adults often have earlier, lighter and more fragmented sleep, but age does not remove the need for safe daytime alertness.

Chronotype adds variation in timing. Genetics contributes, while age, light, work and habits shape how it appears. A late chronotype is not laziness. It becomes a problem when internal time and external obligation collide.

Someone who sleeps midnight to eight on free days but rises at six for work repeatedly crosses between schedules. Social jet lag names that mismatch; it does not prove that every late weekend is disease.

Duration is one dimension within a broader concept of sleep health. Continuity asks whether sleep is repeatedly broken. Timing asks whether it occurs at a suitable biological and social phase. Regularity asks how much the schedule moves from day to day. Satisfaction captures the sleeper's experience. Daytime alertness asks whether wakefulness works. Breathing, movement, architecture and freedom from a disorder add information that time alone cannot provide. The dimensions overlap, and no accepted consumer score combines them into one clinical truth.

This changes how personal need is estimated. A single alarm-free morning is poor evidence after a restricted week. A longer period with adequate opportunity, fairly stable timing and no large accumulated debt is more informative. Daytime function matters, but it is not infallible because self-assessment can adapt. Unintended sleep, repeated emergency caffeine, irritability, concentration failures and drowsy driving are stronger warnings than pride in a short schedule.

Long sleep requires equal care. In population studies, long reported duration often accompanies poor health. That does not establish that sleep caused the illness. Depression, medication, frailty, inflammation and fragmented nights can increase time in bed or reported sleep. Cutting sleep solely because an epidemiological curve rises at the far end would confuse marker with cause.

The social dimension belongs inside the model. Shift work, multiple jobs, caregiving, overcrowding, heat, noise and unsafe housing distribute opportunity unequally. A person may understand every sleep rule and lack the power to follow it. Personal routines can improve the part of the system they reach. They should not rename an employment, housing or health problem as weak discipline.

The practical target is therefore a pattern: enough opportunity, workable

timing, reasonable regularity, tolerable continuity and wakefulness that remains alert and safe. Persistent difficulty under those conditions is useful information. Individual variation widens the range. It does not abolish the need or reduce health to one number.

The Day Builds the Night

A good night begins long before the lights go out. This is unwelcome news for anyone selling a product that can be placed on a bedside table.

Light is the strongest environmental time cue for the human circadian system. Light after waking often supports alignment with the local day and, when the clock is delayed, can help shift timing earlier. Bright evening and early-night light tends to push timing later. The direction and size depend on intensity, duration, spectrum and the person's biological phase. Outdoor light is usually stronger than ordinary indoor lighting, while a brightly lit room or screen close to the face can remain relevant at night.

Regular wake timing can give the clock a repeated reference, anchor morning light and provide a predictable period for sleep pressure to build. It is a powerful lever for many people, not a universal master rule. A rotating night worker, a parent feeding an infant or someone with a circadian disorder may need a deliberately different plan. Bedtime is less obedient than rise time. Entering bed because an ideal schedule says so, while alert, can create an hour of rehearsal in wakefulness.

Activity changes the day-night transition through several routes. Regular exercise generally supports sleep and health. The evidence does not justify banning all evening training. Timing, intensity and individual response matter. A demanding session close to bed may leave one person hot and activated while another sleeps well. The relevant test is the repeated effect on timing, arousal and opportunity.

Caffeine modifies the perception and expression of sleep pressure. Dose matters as much as the clock. In a 2025 randomised crossover study of twenty-three healthy men with moderate habitual use, 100 milligrams taken four hours before bed did not produce a significant group-level effect, while

a single 400-milligram dose altered sleep when taken within twelve hours and increased fragmentation when taken within eight. The study does not create a safe universal cut-off: it included one narrow population and tested fixed doses. It does show why “one coffee” and “after noon” are crude units. Work backwards from dose, timing, pregnancy, medication, sensitivity and the outcome you care about.

Naps spend pressure. A short early-afternoon nap can improve alertness with limited grogginess for many adults. Longer naps may provide more recovery and increase the chance of sleep inertia. Late naps can make night sleep harder. The same nap may be intelligent after acute loss and counterproductive when chronic insomnia is being maintained by low pressure and excessive time in bed.

Screens act through more than colour. Light can shift the clock or reduce evening sleepiness, but content can prolong wakefulness and raise arousal. A dim amber phone still consumes time. The useful question is whether the device is changing light exposure, extending the day, carrying work into bed or doing several at once.

Temperature, food, nicotine, alcohol, medicines, pain and emotion also enter the system. A comfortably cool, dark and quiet room usually helps. A warm bath or shower may support later heat loss. Heavy alcohol can sedate early while damaging later organisation. There is no ideal room number or perfect meal time that outranks comfort, illness, culture and personal response.

A routine succeeds when it removes repeated conflicts and protects opportunity. It fails when it becomes a test whose slightest breach predicts disaster. The strongest evening is often ordinary: the active day has ended, light is lower, the coming schedule is credible, and the bed is no longer a workplace.

Trying Hard Can Keep You Awake

Most actions reward effort. A heavy door opens when pushed harder. A difficult paragraph yields to concentration. Sleep reverses that rule. Effort is wakefulness applied to a state that requires reduced monitoring and control.

An acute bad night is common. Pain, stress, noise, illness, travel, excitement or hot flushes can delay or fragment sleep, and the problem may resolve when the cause passes. Chronic insomnia disorder is different. It involves persistent difficulty initiating or maintaining sleep despite adequate opportunity, with distress or daytime impairment. It can begin with a sound reason for wakefulness and continue after the original disruption has weakened.

One useful account separates predisposing, precipitating and perpetuating factors. A person may begin with a sensitive arousal system or late chronotype. A job crisis precipitates poor nights. In response, bedtime moves earlier, rising moves later, naps multiply, activity shrinks, clocks are checked and every awakening becomes evidence of tomorrow's damage. These responses are understandable. Together they can reduce pressure, destabilise timing and teach the brain that bed predicts effort and threat.

Generic sleep hygiene has limits because it mostly removes obstacles. A dark room, sensible caffeine timing and a workable schedule do not necessarily reverse a learned association between bed and wakeful struggle. The 2023 European insomnia guideline recommends cognitive behavioural therapy for insomnia, CBT-I, as first-line treatment for adults across ages, including people with comorbid conditions. Properly delivered digital CBT-I can widen access. Sleep-hygiene advice alone is not an equivalent treatment.

CBT-I combines methods that target the maintaining mechanisms. Stimulus control rebuilds the association between bed and sleep by reducing prolonged wakeful problem-solving there, using the bed for sleep and rising at a consistent planned time. Time-in-bed restriction or compression temporarily brings opportunity closer to estimated sleep, then expands it as

continuity improves. Cognitive work addresses catastrophic predictions, rigid rules and monitoring. Relaxation and education support the programme without pretending that calmness can be commanded either.

The name “sleep restriction” invites misuse. It is a structured treatment, not a challenge to survive on less sleep. Sleepiness can increase early, and the method may need adaptation or supervision for bipolar disorder, seizure disorders, untreated sleep apnoea, high fall risk, pregnancy, severe medical illness or safety-critical work. A paragraph can explain the mechanism without prescribing an individual window.

Medicines are not one thing, and sedation is not proof of uselessness. Prescription treatment can be reasonable when CBT-I is unavailable, insufficient or unsuitable, or when rapid relief matters, provided the choice reflects diagnosis, benefits, adverse effects and intended duration. A 2026 AASM guideline, based on low-certainty evidence, suggests CBT-I alone over routine combination treatment and combination treatment over medication alone. Those conditional recommendations are not a ban. They mean a sleeping tablet should not displace the treatment that changes the maintaining system. Availability without prescription does not establish effectiveness for chronic insomnia: antihistamines can leave residual effects, while melatonin is chiefly a timing signal rather than a universal treatment for the disorder. The useful question is what problem a product is meant to treat, not whether it sounds natural or sedating.

The central change is a reduction in emergency. A poor night raises pressure for the next one. Extending the morning, napping repeatedly and entering bed far too early can blunt that correction. Clock-checking turns normal awakenings into calculations. Effort raises arousal. Treatment changes the behaviours and predictions that keep wakefulness likely.

This does not make insomnia imaginary. Conditioned arousal changes attention, expectation and physiology. Nor does it mean every persistent complaint is insomnia. Pain, depression, medicines, menopause, breathing disorders, restless legs and circadian disorders can coexist.

Menopause-specific CBT may help sleep problems linked with vasomotor

symptoms, but it does not replace assessment of other causes or treatment choices for the symptoms themselves.

The lesson is precise. When a sleep problem is being maintained by the struggle to control sleep, more struggle will strengthen it. The task is to restore the conditions under which sleep can occur without examination.

Different Failures Need Different Responses

The two-process model explains much of normal timing. The wider gated model explains why one list of habits cannot repair every broken night. Similar mornings can be produced by different failures.

Obstructive sleep apnoea is the clearest example. During sleep, the upper airway repeatedly narrows or closes, reducing or stopping airflow until the brain partly wakes the person and restores breathing. The sleeper may remember none of the arousals. A bed partner may hear loud snoring, pauses, choking or gasping. Morning headache, dry mouth, nocturia, poor concentration and unrefreshing sleep can follow. Higher body weight raises risk, but anatomy, age, sex and muscle control mean that thin people can also have the disorder. A quieter room is not the target. An open airway is.

Restless legs syndrome fails differently. An urge to move, often accompanied by unpleasant sensations, appears or worsens during rest, is strongest in the evening or night and improves temporarily with movement. Iron status, pregnancy, kidney disease, medicines and genetics can contribute. Current clinical guidance makes iron studies and review of aggravating factors part of the first assessment. Iron treatment depends on interpreted results and context, not on self-prescribing a high-dose supplement. Advising the person to lie still and relax can intensify the defining feature.

Narcolepsy destabilises sleep-wake control. Excessive daytime sleepiness is central. Some people also experience cataplexy, a sudden loss of muscle tone often triggered by emotion, alongside sleep paralysis, vivid experiences around sleep transitions and fragmented night sleep. Loss of orexin signalling in narcolepsy type 1 helped reveal how the brain stabilises

wakefulness. This is not ordinary tiredness with a dramatic label.

Many parasomnias arise from incomplete or disordered transitions. Sleepwalking and some night terrors usually emerge from deep non-REM sleep. REM sleep behaviour disorder involves loss of the usual muscle atonia, allowing dreams to be enacted and sometimes causing injury. Occasional childhood events can be benign. New, violent or adult-onset dream enactment deserves assessment because the cause and implications differ.

Circadian rhythm sleep-wake disorders preserve an ability to sleep but place it at the wrong internal time. A person with delayed sleep-wake phase disorder may sleep normally on a late schedule and struggle severely when required to rise early. Shift-work disorder and jet lag also produce conflict between internal time and external demand. The complaint can sound like insomnia at bedtime and sleepiness in the morning, while the mechanism is misalignment.

Other causes sit outside neat sleep-disorder labels. Pain, reflux, asthma, menopausal vasomotor symptoms, pregnancy, medication effects, mood disorders, caring duties and environmental noise can break continuity or remove opportunity. Treating the night without treating the cause may improve a symptom and leave the mechanism intact.

Conditions overlap. A person can have insomnia and sleep apnoea, then retain conditioned wakefulness after breathing improves. Depression can cause and follow disruption without explaining every snore. Medicines can alter breathing, movement, timing or architecture. Consumer devices can raise a question but cannot settle the diagnosis.

The threshold for assessment is a pattern, not one imperfect night. Persistent difficulty despite adequate opportunity, dangerous daytime sleepiness, witnessed breathing pauses, choking or gasping, recurrent irresistible leg symptoms, sudden sleep attacks, cataplexy, injurious behaviour or a major unexplained change all move the problem beyond routine advice.

The conditions that permit and maintain sleep also explain how it fails. When pressure is low, timing is displaced, arousal is learned, opportunity is absent or continuity is repeatedly broken, the same complaint demands a different response. Better routines can align a working system. They cannot open an airway, stabilise narcolepsy, remove a night shift, treat iron deficiency or reverse chronic insomnia by persuasion. Real rest begins with matching the response to the failure.

How It Actually Works

Morning sets internal time

Light reaches the retina before breakfast has done anything useful. Some of it enters the visual system and becomes sight. Some reaches specialised light-sensitive cells that report environmental brightness to the circadian clock. The signal helps anchor the internal day, suppresses melatonin and supports the transition towards wakefulness.

The clock is not reset like a digital watch at one instant. It is nudged by the timing and pattern of light across days. Light in the biological morning generally shifts human timing earlier; light in the late biological evening and early night tends to shift it later. The response depends on where the clock already is. The same lamp can have a different effect on a night worker, a teenager and a traveller who has crossed eight time zones.

Wakefulness is maintained by several interacting brain systems rather than one centre. Networks using orexin, histamine, noradrenaline, dopamine, acetylcholine and other signals help stabilise the waking state. Orexin is especially important as a coordinator. Its loss in narcolepsy type 1 helps explain why wakefulness can collapse and REM features can intrude at the wrong time.

Meanwhile, homeostatic pressure begins to rise. Adenosine participates in the bookkeeping, though no single molecule is a complete gauge. Caffeine occupies adenosine receptors and makes the increasing pressure less legible. The person may feel restored while the account remains open.

The waking day loads the system

By late morning, the two main forces often cooperate. Sleep pressure has not risen far, while the circadian system strongly promotes wakefulness. Attention, body temperature and many physiological functions follow their own daily rhythms, so performance does not stay flat from breakfast to bed.

After lunch, many people experience a dip in alertness even when the meal is modest. The clock contributes to this afternoon valley. A brief nap can improve alertness while lowering the chance of reaching deeper sleep, but human sleep does not obey a stopwatch. A longer nap can be more restorative after acute loss and more likely to produce sleep inertia on waking. It also subtracts more pressure from the coming night.

Activity, social contact and light reinforce the waking state. Exercise changes temperature, autonomic activity and later sleep propensity, and its long-term effects are generally favourable. Meal timing, activity and work schedules can influence rhythms outside the brain, while some medicines alter sleep or circadian timing. The master clock coordinates the system, but peripheral tissues also keep time. This is one reason circadian disruption is larger than feeling sleepy at the wrong hour.

Caffeine taken now may still matter at bedtime. Clearance varies markedly between people. Pregnancy can slow it; smoking can speed it; medicines and genetics alter it. The dose is often hidden by the vessel. A coffee-shop drink may contain far more caffeine than a small home cup, and an energy product may be consumed quickly. A recent crossover experiment also showed why dose and timing must be read together: 400 milligrams affected later sleep far earlier in the day than 100 milligrams did in the tested group. “One coffee” is not a unit of pharmacology.

Evening creates a second wind

As wakefulness continues, pressure rises. Yet many people feel more alert in the early evening than they did at 3 p.m. The circadian system is producing a strong wake signal that opposes the accumulated pressure. Sleep researchers sometimes call this the wake maintenance zone. It helps keep sleep consolidated into the night instead of letting it begin whenever pressure first becomes uncomfortable.

This second wind can be mistaken for recovered energy. It is better understood as timing. If the person stays awake into the biological night, circadian wake promotion falls while pressure remains high. Sleep becomes easier. If bright light, exciting work or travel pushes the clock later, that opening moves too.

Melatonin begins to rise in dim light before habitual sleep for many people. It is a useful marker of internal night and a signal to several tissues. The rise does not guarantee sleep. A person can produce melatonin while remaining awake through stress, pain, movement or choice; light can also weaken or delay the signal. Taking melatonin at an arbitrary dose and hour may therefore add a signal without solving the timing problem. Its best-supported uses are tied to circadian timing and particular clinical contexts, not a general promise of stronger sleep.

Body temperature also changes. Core temperature tends to decline across the evening, aided by heat loss through the skin. Warm hands and feet can therefore accompany a falling core temperature. A warm bath before bed may seem contrary to the advice to sleep cool, but the warming can increase peripheral blood flow and support heat loss afterwards. The bedroom then needs to permit comfort, not meet a number printed by a mattress company.

Sleep begins by reducing access

Falling asleep is not the brain shutting down in one movement.

Wake-promoting systems lose dominance, sleep-promoting regions inhibit them, and the brain becomes less responsive to the outside world. N1 is the unstable crossing. Thoughts loosen, images appear, muscles may jerk, and a person awakened quickly may insist that no sleep occurred.

N2 makes the state more secure. Eye movements stop, muscle activity falls, and the electroencephalogram, EEG, shows spindles and K-complexes. These patterns are not decorative labels. Spindles are linked to sensory filtering and memory processes. K-complexes can arise spontaneously or follow an external stimulus, marking a large cortical response that does not necessarily become full awakening.

Sleep therefore reduces responsiveness rather than abolishing it. Regular and predictable input is often less disruptive than sudden change, although the effect depends on intensity, meaning, stage and the sleeper. Earplugs or steady background sound can reduce some disturbances. They do not switch the sensory system off, and an abrupt or personally significant event may still trigger arousal.

As pressure remains high, sleep deepens into N3. Large slow waves reflect broad coordination across cortical networks. Heart rate and blood pressure tend to fall, breathing becomes more regular, and waking the sleeper becomes harder. Growth hormone secretion is associated with early sleep, especially slow-wave sleep, though repair cannot be assigned to one hormone or stage.

Deep sleep is concentrated early because homeostatic pressure is highest at the start. After deprivation, the brain often protects slow-wave activity and intensity. This rebound is evidence that sleep is regulated rather than passively accumulated. It is also why a first cycle after a long day can feel like falling through the floor.

The night changes its composition

Later in the first cycle, with timing that varies widely, sleep moves towards REM. The EEG becomes more wake-like, the eyes move rapidly, and skeletal muscles are largely paralysed by brainstem mechanisms. Breathing and heart rate become less regular. Thermoregulation is reduced. Dreams can become vivid, but dreaming is not exclusive to REM.

The atonia is protective. In REM sleep behaviour disorder, that protection fails and dream enactment can emerge. At the other edge, REM atonia can persist briefly into wakefulness, producing sleep paralysis: awareness returns while voluntary movement does not. The experience can be terrifying and can include vivid perceptions as dream imagery and waking sensation overlap. Isolated episodes can occur without narcolepsy and are usually brief; recurrent events with severe sleepiness or other symptoms may require assessment.

The cycle then returns through lighter sleep and may descend again. Each pass differs. Early cycles contain more N3. Later cycles contain longer REM periods. Brief awakenings occur at transitions and are often forgotten. A person who recalls waking twice may have woken more often without forming a durable memory.

External events keep testing the system. A partner moves. A siren passes. The bladder fills. The airway narrows. Pain changes. Temperature rises under the duvet. Most events are handled without full awakening. Some produce an arousal lasting seconds, enough to alter EEG and physiology but too short to enter memory. Repeated arousals can make a long night unrefreshing.

Breathing deserves special attention. When awake, muscles help keep the upper airway open. During sleep, muscle tone falls. In a susceptible airway, soft tissue narrows the passage, airflow falls, oxygen may drop, carbon dioxide may rise, and the brain triggers an arousal to restore tone. The cycle can repeat many times. The person may believe they slept continuously because the protective awakenings were too brief to remember.

Alcohol can worsen this arrangement. Its early sedative effect may hasten sleep, while reduced REM, airway effects in susceptible people and other later consequences can reduce the quality of what follows. The drink solves the complaint the person can observe, waiting to fall asleep, and can worsen events they cannot observe.

A bad night can train the next one

Consider an illustrative sequence assembled from common insomnia mechanisms, not a reported patient. A stressful event delays sleep. The following day feels impaired, so bedtime is moved earlier. At night the person checks the clock, calculates the remaining hours and monitors every sign of alertness. The bed now contains a task: prove that sleep is happening before tomorrow is damaged.

The monitoring changes the state being monitored. Attention turns towards heartbeat, thoughts, room sounds and time. Each sign of wakefulness gains meaning. The person remains in bed because leaving feels dangerous, yet the extended wakefulness strengthens the association between bed and alert effort. A nap the next day reduces pressure. Another early bedtime creates another long opportunity for wakefulness. The original stress may pass while the loop continues.

Normal sleep is less conscious than this. People misjudge sleep latency because N1 and brief N2 can feel like wakefulness, and they often forget short periods of sleep between remembered awakenings. In insomnia, this gap between experienced and measured sleep can widen without making the distress false. The experience of lying awake is the problem the person lives, even when a laboratory finds more sleep than expected.

CBT-I alters the operating conditions. A fixed rise time restores a reliable anchor. Stimulus control removes prolonged wakeful struggle from the bed. A carefully limited sleep window concentrates pressure and continuity. Cognitive work reduces the catastrophic meaning assigned to one night. Improvement often arrives unevenly because the treatment is changing a learned prediction, not administering instant unconsciousness. The first

target is a bed that no longer announces an examination.

Dawn promotes wakefulness before the alarm

Towards morning, sleep pressure has fallen and the circadian system is moving towards wake promotion. REM periods are longer, deep N3 is less common, and awakenings become easier. Cortisol rises before habitual waking as part of a broader daily rhythm, but calling it a stress surge misses its normal role in preparing the body for the active phase.

An alarm can cut through any stage. Waking from deeper sleep, at an adverse circadian phase or after restriction increases the chance of sleep inertia. The person is awake in the legal sense and not yet fully operational. Inertia usually fades, but it can last longer after severe loss, night work or particular disorders. Repeated snoozing adds fragmented sleep and repeated awakenings; it does not guarantee a restorative extra cycle.

Morning light then begins the next adjustment. The clock, pressure and schedule begin interacting again. The strength of the system lies in repetition. One late meal, one bright screen or one short night rarely destroys sleep. A pattern does more than an isolated event because it keeps teaching the clock, altering pressure and changing what the bed predicts.

What happens across a week

A weekday schedule often creates a hidden experiment. The alarm fixes wake time while work, care and entertainment delay bed. Pressure rises across successive short nights. The person compensates with caffeine, sleeps later at the weekend and feels better. Recovery sleep is useful, and consensus work supports the value of catch-up when weekday duration has been inadequate. A large change in wake time can also shift light exposure and reduce pressure for Sunday night. Monday may therefore begin with less debt than Friday and more circadian displacement than Saturday.

Controlled studies of weekend recovery show why the argument cannot be settled by slogans. Extra sleep can improve alertness and repay part of an

acute deficit. It does not reliably erase every metabolic or cognitive effect of repeated restriction, especially when the pattern resumes. The correct conclusion is neither “sleep debt can never be repaid” nor “one lie-in fixes the week”. Recovery helps. Prevention is more complete.

Shift work makes the conflict harder because the schedule itself rotates. Night workers must perform near the circadian low and try to sleep when light and the clock promote wakefulness. Strategic light, darkness, naps, caffeine timing and schedule design can reduce harm, but a person cannot become indifferent to time through practice. Some adapt better than others; frequent rotation may prevent stable adaptation altogether.

Jet lag is the temporary version. The traveller's clock remains tied to the departure zone while the destination demands a new pattern. Eastward travel often requires an advance, sleeping and waking earlier by internal time, which many people find harder than delaying westward. Light timed to the new schedule is the main resetting cue. Melatonin can help in some situations, but timing determines whether it shifts the clock in the intended direction.

The clinic reconstructs the hidden night

A sleep complaint begins with a history, not a stage score. The clinician asks when sleep is attempted, when it occurs, how it differs across workdays and free days, what the daytime consequences are, which medicines and substances are involved, and what another person has witnessed. A two-week diary can reveal delayed timing, variable opportunity or long periods in bed. Actigraphy can add an objective estimate of movement and rest across ordinary life. Neither tool explains every awakening.

The next test depends on the suspected mechanism. Chronic insomnia can often be diagnosed from symptoms and context without an overnight laboratory study. Suspected obstructive sleep apnoea may be investigated with a home respiratory study or full polysomnography, depending on complexity and local guidance. Airflow, breathing effort and oxygen signals

reveal events that a wrist device cannot. The apnoea-hypopnoea index counts events per hour, but symptoms, oxygen burden, sleep position, stage and other findings affect interpretation. A threshold is a classification aid, not the whole patient.

Restless legs syndrome is identified mainly through its characteristic urge, timing, rest trigger and relief with movement; blood tests may look for iron deficiency and other contributors. Narcolepsy assessment may combine an overnight study with a multiple sleep latency test, which measures how quickly sleep begins during scheduled daytime opportunities and whether REM appears unusually early. Parasomnias may require video and EEG when events are dangerous, unusual or hard to distinguish from seizures.

The method is differential rather than ceremonial. A person can report “I sleep badly” because sleep is too short, mistimed, fragmented by breathing, interrupted by movement, destabilised by neurology, conditioned by insomnia or displaced by pain and mood. The clinic's job is to find which hidden night produced the visible day.

The science learned to see sleep

For most of history, sleep could be observed only from outside and reported from memory. EEG changed that. In the 1930s, Alfred Loomis and colleagues described recurring electrical patterns across sleep. Aserinsky and Kleitman's 1953 REM report added eye movements and a distinct state. William Dement and Kleitman then helped establish the cyclic organisation of the night.

Modern polysomnography records several signals at once: brain activity, eye movements, muscle tone, airflow, breathing effort, oxygen saturation, heart rhythm and often leg movement. The combination matters. An EEG pattern can stage sleep; belts and airflow sensors show whether breathing continues; oxygen reveals physiological consequence; video can capture behaviour.

Wearables work with a smaller view. Wrist movement and optical pulse signals estimate sleep and wake, then algorithms infer stages. Reviews and

device comparisons continue to find substantial variation across products, outcomes and people. Sleep-wake estimates are generally more dependable than exact stage minutes, and proprietary updates can alter performance without changing the hardware. The output can reveal patterns and remains weak as a verdict. A clinical diagnosis requires the signals and judgement suited to the suspected disorder.

How we know

Sleep science combines controlled restriction, circadian protocols, polysomnography, field measurement, animal experiments, clinical trials and long-term cohorts. Each answers a different question. Restriction experiments support short-term causal claims but do not reproduce every ordinary poor night. Cohorts reveal longer patterns while leaving illness, occupation and social conditions entangled. Animal work can expose mechanisms inaccessible in humans, and translation remains a separate test.

Sleep stages are operational categories built from brain activity, eye movement and muscle tone. They are useful maps with imperfect borders. Wrist devices infer them from fewer signals and do not become polysomnographs through numerical precision.

Memory effects differ by task and sleep feature. Mouse clearance experiments in 2013 and 2024 reached opposing conclusions with different tracer methods, while 2025 work linked NREM vascular rhythms to fluid movement. A 2026 human crossover study found biomarker patterns consistent with sleep-active clearance but relied on an investigational device and mathematical model. That is evidence, not closure. Long sleep may mark illness rather than cause it, and large studies often use self-reported duration.

The secure core is narrower: sleep and circadian timing are regulated; the night contains recurring physiological states; repeated restriction impairs waking function; and disorders can fragment, displace or destabilise sleep in ways that hours in bed will not reveal.

What People Get Wrong

“Eight hours is the whole target”

Eight is a useful round number, not a biological verdict and not a complete definition of sleep health. Expert consensus advises adults to obtain at least seven hours regularly, while recognising individual variation and circumstances in which more is appropriate. Age, pregnancy, illness, previous loss and daily demand can alter need.

The correction does not justify self-deception. Natural short sleepers exist, but they are rare. Identified inherited phenotypes come from family and laboratory research, not from asking whether five hours feels normal. Familiarity with restriction is weak evidence because subjective sleepiness can flatten while objective lapses continue to accumulate.

Duration also cannot identify continuity, timing, regularity, breathing or daytime alertness. Seven hours repeatedly broken by airway obstruction is not equivalent to seven consolidated hours. Eight hours at a severely misaligned biological time may remain fragile. A person can sleep an acceptable duration and still require assessment for persistent sleepiness.

A fixed number became attractive because it is easy to count and sell. The dimensions that often explain a poor night are messier: when the sleep occurred, how often it broke, whether the airway remained open and whether the schedule repeated. Precision on duration can conceal ignorance elsewhere.

The better question is whether the pattern regularly permits alert, safe wakefulness without repeated unplanned sleep, escalating caffeine or large recovery swings. The number is a screening clue. It is neither the whole target nor the diagnosis.

“If I cannot sleep, I should stay in bed and try harder”

This feels sensible because leaving bed appears to surrender. In chronic insomnia, it can strengthen the problem. Long periods awake in bed pair the bedroom with monitoring, frustration and effort. Going to bed earlier can spread the same amount of sleep across a larger opportunity, lowering pressure at the intended time and creating more wakefulness to notice.

Stimulus control within CBT-I reverses that learning. The bed is reserved for sleep rather than prolonged wakeful problem-solving. The person goes when sleepy, follows a planned rise time and leaves the bed during sustained struggle until sleepiness returns. The exact plan must fit mobility, safety and clinical context.

The principle is not punishment, and every brief awakening does not require evacuation. It interrupts the association that matters. A bed should predict reduced vigilance. If it has become an examination hall, more time sitting the examination is poor preparation. A rigid twenty-minute rule can make matters worse by encouraging clock-checking. The aim is to stop rehearsing wakeful effort, not create another performance metric.

People with bipolar disorder, epilepsy, falls risk or severe daytime sleepiness may need a modified plan rather than self-directed restriction of time in bed. The mechanism remains useful; implementation becomes clinical when the cost of additional sleepiness is high.

“Alcohol helps me sleep”

Alcohol can shorten the wait for sleep at higher doses, which makes the claim feel proven. The measurement stops too early. Controlled studies and recent review evidence show delayed REM onset and reduced REM duration even at lower doses, with stronger effects as dose rises. Effects on total sleep time, efficiency and wakefulness after sleep onset are less consistent. Urination, temperature changes and poorer breathing in susceptible people can add to the cost.

Effects vary with dose, timing, tolerance, sex, body size, medication and the presence of sleep apnoea. A single drink does not guarantee a ruined night, and one precise universal cut-off would be false confidence. The stable distinction is between becoming less responsive and receiving well-organised sleep.

The 2025 meta-analysis found a clearer dose-related effect on REM than on total sleep time, sleep efficiency or wake after sleep onset, where estimates remained uncertain. That is a warning against converting one plausible mechanism into a complete prediction of the night.

Using alcohol as a sleep tool also treats a visible symptom, waiting to fall asleep, while leaving timing, pressure, arousal or diagnosis untouched. In a vulnerable airway, alcohol can worsen snoring and obstructive events. The nightcap may solve the only part of the night the drinker watches.

“Blue light is the whole screen problem”

Short-wavelength light can influence the circadian system, and sufficiently bright evening screens can suppress melatonin and delay timing. Colour is only one variable. Brightness, duration, distance, timing and biological phase all affect the signal.

Screens also consume time and attention. A dim amber phone can delay bedtime, deliver work conflict, provoke comparison or keep novelty arriving. A night setting may reduce one part of the light exposure while allowing the behaviour to continue for another hour.

Lower brightness and warmer settings can help, especially for people with late timing or strong sensitivity. They are not permission slips. Ask whether the device is shifting the clock, extending wakefulness, raising arousal or doing several at once. Laboratory e-reader studies used prolonged controlled exposure and do not prove that every brief glance has the same effect. Screen light is biologically relevant; “blue” should not become a magic word that hides dose and behaviour.

The same light can shift timing differently according to biological phase.

Morning light often advances the clock; late-evening light often delays it. Bright-light treatment and deliberate darkness can therefore be useful tools, but only when the timing problem has been identified.

“I can repay sleep debt in one weekend”

Extra sleep after restriction is useful. Pressure rises during loss, and recovery sleep can improve alertness and restore some performance. Consensus work supports catch-up sleep when weekday duration has been insufficient. The myth lies in treating one weekend as a complete and instant settlement.

Controlled studies show that some cognitive or metabolic effects can persist after one or two longer nights, particularly when restriction has been repeated and then resumes. Sleeping much later can also move circadian timing, making the next early start harder. The size of each effect depends on the person and schedule.

Refusing a needed lie-in would be the wrong lesson. Recovery is sensible. Use it while recognising the trade-off. A weekend can reduce an acute deficit; it cannot make a chronically inadequate weekday schedule harmless. When feasible, combining earlier bedtimes with a modestly later morning may recover duration while moving the clock less. The most complete repayment plan stops taking out the same loan.

“Snoring is harmless”

Simple snoring can occur without obstructive sleep apnoea. The sound alone is not a diagnosis. The dangerous mistake is to treat every snore as comic when it is accompanied by breathing pauses, choking, gasping, unrefreshing sleep, morning headaches or excessive daytime sleepiness.

In obstructive sleep apnoea, the upper airway repeatedly narrows or closes. The brain restores breathing through arousal, often without a remembered awakening. Time in bed can look adequate while airflow, oxygen and continuity are repeatedly disturbed. Higher body weight raises risk, but anatomy and other factors mean the disorder is not confined to one body

type.

A phone recording or tracker may raise suspicion. It cannot rule the disorder in or out. Assessment uses symptoms, witnessed events and respiratory testing or polysomnography according to context. The correction matters because the person often cannot observe the event. A problem hidden inside sleep may be first visible to somebody sharing the room.

“My tracker knows how I slept”

A wrist device knows movement, pulse-derived signals and whatever else its sensors capture. It does not ordinarily measure brain waves, eye movements, chin muscle tone, airflow, breathing effort and oxygen together. Its stages are modelled estimates.

That does not make the device useless. It can reveal bedtimes, rise times, broad duration patterns and associations with travel, alcohol or training. A 2026 rapid review of twenty-nine ambulatory studies found moderate accuracy for total sleep time and time in bed, with lower precision for sleep efficiency, wake after sleep onset and stage classification. REM and deep-sleep estimates were especially unreliable, and clinical populations were rarely represented. Performance still varies across brands, metrics, people and nights, and an algorithm update can change the answer without changing the sleeper.

The score becomes harmful when it outranks function. Someone feels restored, sees a poor rating and becomes anxious; the anxiety then damages the next night. Another person feels persistently sleepy and accepts a reassuring score. Use the device as a pattern detector, not an authority. Persistent sleepiness, insomnia, breathing symptoms or unusual behaviour require history and appropriate testing. Orthosomnia describes the trap in which pursuit of perfect data begins to injure the sleep being measured.

Use It

Separate opportunity from ability

Begin by asking whether sleep had a fair chance. Sleep opportunity is the interval protected for sleep, not the number displayed the next morning. It must include ordinary latency, brief awakenings and the transition out of bed. A schedule offering six hours cannot be repaired by efficiency.

This separates shortage from inability. Someone working late and rising early may need more protected time. Someone spending ten anxious hours in bed and sleeping six may need a different approach because increasing opportunity has already failed. The same complaint, “I did not get enough sleep”, points in opposite directions.

Examine a week rather than one night. When did work, travel, care or entertainment end? Were there enough nights with credible opportunity? Did recovery depend on large weekend shifts? The first intervention may be a boundary around time, not a supplement or new mattress.

Stabilise timing without worshipping sameness

Circadian systems learn from repeated light and behaviour. A reasonably consistent wake period and useful light after waking often provide a strong anchor. Evening light, late work and repeated shifts in schedule can move the biological night later. Regularity matters because the clock cannot align with a target that changes continually.

Consistency is a tool rather than a moral code. A late chronotype may need a later stable schedule. A night worker may need carefully timed light and darkness. Parents, carers and people with fluctuating illness may be unable to preserve a narrow wake time. The objective is to reduce avoidable movement while respecting the life that exists.

Judge timing by pattern. Do you sleep more easily and wake more naturally on free days at a different phase? Does an early obligation produce insomnia at night and sleepiness in the morning? Does travel or shift

rotation explain the change? A timing problem often looks like weak discipline until the clock is allowed to speak.

Trace the failing condition

Use four questions after a repeated poor night. Was homeostatic pressure too low because of a late nap, little wake time or excessive time in bed? Was circadian timing displaced by light, travel, chronotype or shifts? Was arousal maintained by pain, threat, work, excitement or the effort to sleep? Was continuity broken by breathing, movement, temperature, hot flushes, medication, care or noise?

These questions do not diagnose. They prevent category errors. Earlier bedtime helps a compressed schedule and fails against delayed timing. Relaxation may lower arousal and cannot open an airway. More time in bed can repair shortage and perpetuate some insomnia. Earplugs can reduce noise and cannot treat restless legs.

Choose the response that reaches the suspected mechanism, then watch whether the predicted pattern changes. When the answer remains unclear, persistent or dangerous, clinical assessment becomes the next experiment. Owning a category is more useful than collecting twenty unrelated tips.

Test one lever at a time

Many sleep rules are compressed averages. Caffeine effects depend on dose, timing and metabolism. Evening exercise depends on intensity and individual response. Naps change both alertness and later pressure. Screen use combines light, time and content. Turn the uncertain advice into a proportionate personal test.

Keep the main schedule reasonably steady. Change one material variable for long enough to see a pattern. Record only what can inform a decision: opportunity, estimated latency, remembered awakenings, rise time, caffeine dose and timing, and daytime alertness. A paper diary often supplies enough evidence. Change six variables at once and the most expensive purchase will claim the credit.

Stop when the experiment has answered the question. Someone who sleeps well does not need a permanent laboratory. Someone with chronic insomnia may need CBT-I rather than more nightly observation. Self-tracking has failed when it increases vigilance or makes a normal fluctuation feel dangerous.

Judge the night in daylight

Sleep is private and mostly forgotten. Its purpose is visible during wakefulness. Ask whether you remain alert in meetings, while reading, during training and on routine journeys. Notice unintended dozing, repeated concentration lapses, irritability, reliance on emergency caffeine and whether another person reports breathing pauses or unusual behaviour.

Daytime function corrects both pride and panic. Someone can celebrate five-hour nights while fighting sleep and making avoidable errors. Someone else can feel restored after a night the tracker disliked. Function is still imperfect because people adapt subjectively and many illnesses produce fatigue without sleepiness. It belongs beside duration and pattern, not underneath a score.

Safety outranks interpretation. Struggling to keep the eyes open while driving or operating machinery is not a moment for open windows, loud music or a breathing exercise. Stop the hazardous activity and arrange a safe opportunity to sleep. Recurrent dangerous sleepiness needs assessment because voluntary restriction is only one possible cause.

Build a routine that survives real life

A useful routine reduces decisions and conflict. It may include lower light, a clear end to work, preparation for morning, a quiet activity, medication taken as prescribed and a room that permits sleep. Its value comes from repetition and fit, not ceremonial length.

Keep a minimum version that survives travel, deadlines and family events. Protect enough opportunity, local light after waking, sensible caffeine timing and a credible end to the active day. A robust routine bends. A brittle one

converts a late dinner into a week of abandonment.

Households are part of the system. Partners can differ in chronotype, temperature, alarms and movement. Children and relatives can interrupt opportunity. Snoring can be another person's first evidence of disease. Solutions may include separate duvets, earplugs, changed alarm methods, redistribution of night care, medical assessment or occasional separate sleep. Identical thermoregulation is not a test of intimacy.

The limits

A general model cannot choose treatment for an individual. Persistent insomnia, suspected sleep apnoea, repeated irresistible urges to move the legs, episodes of sudden sleep, cataplexy, injurious sleep behaviour, severe mood disturbance and dangerous daytime sleepiness require professional assessment. Pregnancy, bipolar disorder, neurological disease, medication changes, falls risk and safety-critical work can alter what is appropriate.

Opportunity is also social. A rotating nurse, an infant's parent, a worker holding two jobs and a tenant beside a nightclub do not face the same choice set as someone comparing blackout blinds. Sleep is biological, while time, quiet and safety are distributed through work, money, housing and care. Advice that ignores those constraints converts an institutional problem into personal blame.

Sleep cannot carry every health ambition. It supports waking function and physiology. It does not cancel smoking, inactivity, poor diet, untreated disease or chronic stress. Persistent fatigue can also arise from causes outside sleep. A good model tells you when to stop adding sleep advice and widen the investigation.

The one thing to keep

Keep the conditions.

Sleep is not a button and not a score. It becomes possible when pressure, circadian timing, opportunity and arousal line up, then remains useful only if breathing, movement, temperature and the environment allow continuity.

Most sound routines work by improving one of those conditions. Good treatment identifies the condition that routines cannot reach.

This changes the question after a poor night. Instead of “Why did I fail?”, ask what the pattern predicts. A late nap points towards pressure. Easy sleep on free days at a later time points towards circadian mismatch. A bed full of calculations points towards learned arousal. Witnessed pauses point towards breathing. The answers are provisional, but they lead in different directions.

It also changes control. Direct control is weak because nobody can order the next stage or guarantee unconsciousness by a deadline. Indirect control is considerable: time, light, activity, opportunity, associations, household design and access to care can all be changed. The skill lies in using that control without turning sleep into a performance.

The final judgement arrives in daylight. Real rest is present when wakefulness is available for the life that follows: alert enough to notice risk, steady enough to think, and safe enough to act. The missing hours matter because they alter the person who must use the remaining ones.

Terms

Sleep health. A multidimensional description across duration, continuity, timing, regularity, satisfaction, alertness, architecture and disruptive disorders. No single score captures every dimension.

Sleep architecture. The pattern and distribution of sleep stages across a night. Duration can remain unchanged while continuity, timing and stage composition differ sharply.

Circadian rhythm. A biological cycle of roughly twenty-four hours. Sleep timing, alertness, temperature and hormone release are among the many functions shaped by circadian time.

Zeitgeber. German for time-giver: an environmental cue that adjusts biological clocks. Light is the strongest human example; meals, activity and

social schedules contribute. Timing determines the direction.

Suprachiasmatic nucleus. The brain's main circadian coordinator, above the optic chiasm. Retinal light signals align internal time with day and night, and the SCN coordinates rhythms elsewhere.

Phase shift. A movement in biological timing. An advance moves rhythms earlier; a delay moves them later. Light or melatonin can produce either direction according to when it is applied.

Melatonin. A hormone released during biological night. It signals darkness and can shift timing when used appropriately. It is not a universal sedative switch.

Homeostatic sleep pressure. The drive to sleep that generally strengthens across wakefulness and diminishes as sleep proceeds. Naps and previous sleep alter the pressure; caffeine changes part of its felt signal.

Adenosine. A signalling molecule involved in sleep homeostasis. Caffeine blocks adenosine receptors rather than cancelling prior wakefulness. Adenosine is one part of the regulation, not a complete molecular gauge.

Two-process model. The framework in which homeostatic pressure interacts with circadian timing to regulate sleep and wake. It explains naps, second winds, jet lag and tired-but-alert states. It does not describe every cause of insomnia or fragmentation.

Chronotype. A tendency towards earlier or later sleep and activity. Genes contribute, while age, light, work and habits shape its expression. It is a distribution, not two fixed types.

Social jet lag. The recurring mismatch between biological timing and social schedules, often visible as different sleep times on workdays and free days. It names a timing conflict, not a formal disorder by itself.

Sleep opportunity. The interval available for sleep, which exceeds sleep duration because latency and awakenings use part of it. A schedule can protect opportunity directly.

Sleep latency. The time between attempting sleep and falling asleep. Subjective estimates can be inaccurate when light sleep is experienced as wakefulness.

Wake after sleep onset. WASO is the total time awake after first falling asleep and before final waking. It is one measure of continuity and says nothing by itself about the cause.

Sleep efficiency. Estimated sleep time divided by time in bed, expressed as a percentage. It is useful in clinical context and poor as a perfection target. High efficiency can coexist with inadequate duration.

Non-REM sleep. Sleep outside REM, divided into N1, N2 and N3. It ranges from the unstable transition into sleep to deep slow-wave sleep and contains changing brain and body states.

N1. The light, unstable boundary between wakefulness and sleep. Someone awakened from N1 may believe no sleep occurred.

N2. Established non-REM sleep containing spindles and K-complexes. It often occupies more of an adult night than any other stage and is not disposable filler.

N3. Deep slow-wave sleep, weighted towards the earlier night. It reflects high homeostatic pressure and broad coordination of cortical activity. Its amount varies with age, prior sleep and other conditions.

REM sleep. Rapid eye movement sleep, marked by active brain patterns, rapid eye movements and reduced skeletal muscle tone. It becomes more prominent later in the night. Dreaming occurs in REM and non-REM sleep.

Sleep spindle. A brief burst of rhythmic EEG activity characteristic of N2. Spindles are linked to sensory gating and several forms of learning and memory processing.

Arousal. A brief shift towards lighter sleep or wakefulness. Arousals can protect breathing or respond to disturbance. Repeated events may fragment sleep without entering memory, so their cause matters more than the count

alone.

Sleep inertia. Reduced alertness and performance immediately after waking. It can be stronger after deep sleep, circadian misalignment or substantial sleep loss. Being upright does not guarantee full readiness.

Microsleep. A brief involuntary episode of sleep during intended wakefulness. It may last only seconds and creates serious risk during driving or safety-critical work. The sleeper may not recognise it.

Insomnia disorder. A lasting pattern of trouble starting or sustaining sleep when there is enough chance to sleep, together with distress or impaired daytime function. It is more than an occasional bad night and can be maintained by learned arousal after the original trigger has passed.

CBT-I. Cognitive behavioural therapy for insomnia. It combines methods such as stimulus control, time-in-bed adjustment and cognitive work to reverse self-maintaining insomnia processes. It is first-line treatment rather than an expanded list of sleep-hygiene tips.

Obstructive sleep apnoea. Repeated narrowing or closure of the upper airway during sleep, causing disrupted airflow and arousals that may not be remembered. Snoring can be present without apnoea, while witnessed pauses or gasping deserve assessment.

Restless legs syndrome. An urge to move the legs, usually worse at rest and in the evening, often accompanied by uncomfortable sensations and temporary relief through movement.

Narcolepsy and cataplexy. Narcolepsy is a neurological disorder of sleep-wake stability causing excessive daytime sleepiness. Cataplexy is sudden muscle weakness, often triggered by emotion, and occurs in narcolepsy type 1. Other features can include sleep paralysis and fragmented night sleep.

Go Deeper

The broad clinical guide

Meir Kryger, *The Mystery of Sleep: Why a Good Night's Rest Is Vital to a Better, Healthier Life* (Yale University Press, 2017). Kryger is a sleep physician and long-standing editor of a major clinical textbook. This is the widest next step: normal sleep, changing needs across life and the disorders hidden inside an apparently full night. Clinical cases keep the medicine human. Read it when this book has made you suspect that “sleep problem” covers several different systems. Its breadth means that no one mechanism receives textbook depth, which is the correct trade for a first general survey.

It also gives breathing disorders and daytime sleepiness enough weight to stop routine advice becoming a universal answer.

The clock

Till Roenneberg, *Internal Time: Chronotypes, Social Jet Lag, and Why You're So Tired* (Harvard University Press, 2012). Roenneberg explains how biological timing meets work, school, light and alarm clocks. It is the strongest continuation of the rhythm half of this book, especially for readers who have treated owlness or larkness as personality. Some health associations are observational, so keep mechanism separate from correlation. The examples still show with unusual clarity how ordinary schedules can create biological conflict without a formal sleep disorder.

The publication predates several newer association studies, but its clock mechanism and schedule analysis remain useful.

The insomnia treatment

Colin Espie, *Overcoming Insomnia 2nd Edition* (Robinson, 2021). This is a structured self-help presentation of CBT-I by a leading researcher and clinician. It explains stimulus control, time-in-bed adjustment, racing thoughts and the logic of changing learned wakefulness. Use it for persistent insomnia rather than occasional shortage. The methods can increase sleepiness at first and are not a substitute for clinical assessment where breathing disorders, bipolar disorder, seizures, pregnancy, falls or safety-critical work complicate treatment.

Its workbook style asks for repeated practice rather than one clever evening, which suits the disorder's learned component.

The disorders through patients

Guy Leschziner, *The Secret World of Sleep: Journeys Through the Nocturnal Mind* (Simon & Schuster UK, 2020). Leschziner uses clinical cases to show what happens when sleepwalking, narcolepsy, apnoea, hallucinations and other disorders disturb the border between sleep and wake. It is the most narrative recommendation here and the best antidote to the idea that every poor night needs the same routine. The cases illuminate mechanisms rather than estimate how common each disorder is. Read it for medicine made visible through lives, not for self-diagnosis.

Its strength is pattern recognition: the same complaint can emerge from entirely different physiology.

Notes and Sources

The Whole Thing in One Page and Why You Should Care

Sleep deficiency and multidimensional sleep health. The National Heart, Lung, and Blood Institute uses *sleep deficiency* for too little sleep, sleep at the wrong time, poor-quality or incomplete sleep, and sleep disrupted by a disorder. St-Onge and colleagues' 2025 American Heart Association scientific statement frames sleep health as multidimensional, including duration, continuity, timing, regularity, satisfaction or quality, daytime alertness or sleepiness, disturbed sleep and architecture. The manuscript uses this as a reader model, not as a validated universal scoring system.

Performance under restriction. Van Dongen and colleagues assigned forty-eight healthy adults aged twenty-one to thirty-eight to eight, six or four hours in bed for fourteen nights, or total sleep deprivation for three nights. The four-hour and six-hour conditions produced cumulative dose-dependent deficits; subjective sleepiness showed much smaller further increases after the initial response. The result supports the narrow claim that people can underestimate accumulating impairment. It is not converted into an exact prediction for every worker, parent or patient.

Safety. Statements about driving and safety-critical work follow the established effects of sleepiness on vigilance, reaction and involuntary lapses. Individual sensitivity, task, circadian phase and recent sleep modify risk. The advice to stop a hazardous activity when struggling to stay awake is a safety boundary, not a method for diagnosing the cause.

Timing, stages and functions

Pressure and circadian timing. Borbély, Daan, Wirz-Justice and Deboer review the two-process model and its limits. The manuscript distinguishes that formal model from the broader practical framework of pressure, timing, opportunity, arousal and continuity. The latter is a synthesis for general readers, not a claim that sleep medicine recognises four official gates.

Adenosine. Reichert, Deboer and Landolt review adenosine and caffeine in sleep-wake regulation. Adenosine is presented as one contributor to homeostasis rather than a substance that accumulates uniformly as a complete measure of sleep need.

Circadian light and melatonin. National Institute of General Medical Sciences material updated in 2025 supports the general description of circadian clocks and light. Auger and colleagues support timed melatonin and light for selected circadian rhythm sleep-wake disorders. The direction of a phase shift depends on biological timing, which is why no universal dose or clock time is supplied.

Stages and cycles. Conventional adult scoring separates wake, N1, N2, N3 and REM using EEG, eye movement and muscle tone, with respiratory and other channels added in clinical polysomnography. Cycle lengths and stage distribution vary; deep N3 is weighted earlier and REM later. Aserinsky and Kleitman's 1953 report is the historical REM anchor, preceded by Loomis, Harvey and Hobart's human EEG work.

Night truncation. Skorucak and colleagues studied seven days of six-hour sleep opportunity and seven days of ten-hour opportunity in thirty-five healthy adults using a crossover design. Restriction primarily reduced REM sleep, while slow-wave activity and accumulated slow-wave activity changed less. The result supports adaptive prioritisation of slow-wave physiology under that protocol, not a universal claim that delayed bedtime preserves all deep sleep or that every curtailed night loses stages in the same proportions.

Learning, immunity and physiology. Rasch and Born review sleep and memory; Besedovsky, Lange and Haack review sleep-immune interactions. The body claims are bounded to experimental changes and organised physiology. Population associations are not treated as proof that short sleep alone causes a specific disease or that extending sleep prevents it.

Brain clearance. Xie and colleagues' 2013 mouse study reported sleep-related changes in interstitial space and clearance. Miao and

colleagues reported reduced clearance during sleep and anaesthesia in male mice in 2024, using a tracer-departure approach that challenged interpretations based on tracer entry. Hauglund and colleagues linked NREM noradrenaline, vascular and cerebrospinal-fluid oscillations to glymphatic flow in mice in 2025. Dagum and colleagues' 2026 randomised crossover study of thirty-nine humans found morning plasma amyloid beta and tau patterns consistent with sleep-active clearance; the inference depended on an investigational device and compartment model. The studies differ in species, tracer, route and outcome. None establishes a settled human detox mechanism or consumer method for enhancing clearance.

Alcohol. Gardiner and colleagues' 2025 systematic review of controlled studies in healthy adults found changes including delayed REM onset and reduced REM duration, with effects related to dose. The text preserves the distinction between shorter latency or sedation and the architecture and continuity of subsequent sleep.

Need, regularity and routines

Adult duration. The American Academy of Sleep Medicine and Sleep Research Society consensus states that adults should regularly obtain seven or more hours to support optimal health. It is used as a population recommendation rather than a personalised target or upper limit.

Natural short sleep. He and colleagues identified a rare DEC2 variant in a short-sleeping family and examined related effects in model systems. This establishes that unusual inherited short-sleep phenotypes exist. It offers no method for an ordinary restricted sleeper to identify as one through confidence or preference.

Age and chronotype. Carskadon's adolescent account supports the interaction of later timing, continued sleep need and early social schedules. Jones and colleagues found many genetic loci associated with morningness in a large genome-wide analysis, supporting biological contribution without genetic destiny. Wittmann and colleagues introduced *social jet lag* for

mismatch between biological and social timing.

Regularity and catch-up. Sletten and colleagues' 2023 National Sleep Foundation consensus concluded that consistent sleep timing matters for health and performance and that catch-up sleep matters when workday sleep is insufficient. The manuscript does not imply that regularity must override recovery or that observational associations prove a precise causal risk.

Caffeine. Gardiner and colleagues' randomised crossover study tested 100 and 400 milligrams at twelve, eight and four hours before bed in twenty-three healthy men with moderate habitual use. The 100-milligram conditions produced no significant group-level objective or subjective effect; 400 milligrams altered sleep within twelve hours and increased fragmentation within eight. The narrow sample and fixed dosing prevent a universal safe cut-off. The study is used to show dose-timing interaction and mismatch between perception and objective sleep.

Screens. Chang and colleagues' controlled e-reader study found circadian and next-morning effects after repeated prolonged evening exposure. The manuscript narrows the conclusion to light, time displacement and arousal rather than treating every screen exposure as equivalent.

Exercise and passive heating. Stutz and colleagues' review does not support a blanket ban on evening exercise in healthy adults, although vigorous late exercise can affect some people. Haghayegh and colleagues found modest average benefits from pre-bed passive body heating across heterogeneous studies. Both justify personal tests rather than universal rules.

Menopause. NICE guidance recommends considering menopause-specific CBT for sleep problems associated with vasomotor symptoms. The book presents this as one option within assessment and treatment, not as a replacement for discussion of hormone therapy, non-hormonal treatment or other sleep disorders.

Insomnia, disorders and measurement

Insomnia and CBT-I. Riemann and colleagues' 2023 European guideline recommends CBT-I as first-line treatment for adults of any age, including people with comorbid conditions, with evidence-based digital delivery as an access route. Edinger and colleagues' AASM guideline supports multicomponent behavioural and psychological treatment and advises against sleep hygiene alone as the sole treatment. Polysomnography is not routinely required for uncomplicated insomnia but becomes relevant when another disorder is suspected or treatment fails. Buysse and colleagues' 2026 AASM guideline gives conditional, low-certainty recommendations favouring CBT-I plus medication over medication alone and favouring CBT-I alone over routine combination treatment. The manuscript preserves room for shared clinical decisions rather than converting that ranking into a ban.

CBT-I cautions. The book explains treatment components without prescribing a personal sleep window. Time-in-bed restriction can initially increase sleepiness and may require modification or supervision where bipolar disorder, seizures, untreated breathing disorders, falls, pregnancy, severe illness or safety-critical work raise risk.

Obstructive sleep apnoea. NHLBI and NICE sources support the account of recurrent upper-airway obstruction, witnessed pauses or gasping, fragmented sleep, risk variation and respiratory testing. The apnoea-hypopnoea index is not treated as the whole clinical picture.

Restless legs, narcolepsy and parasomnias. Winkelman and colleagues' 2025 AASM guideline makes regular iron studies and review of exacerbating factors a central first step for clinically significant restless legs syndrome; thresholds and treatment choices require clinical interpretation. Narcolepsy descriptions follow current National Institute of Neurological Disorders and Stroke material. Cataplexy is defined as sudden muscle weakness often triggered by emotion. New or injurious dream enactment is presented as a reason for assessment rather than a diagnosis.

Consumer devices. Birrer and colleagues' 2024 review found wide

methodological and device variation in wearable sleep staging. Schyvens and colleagues' 2025 comparison of six wrist-worn devices against polysomnography provides a head-to-head example. Landvatter and colleagues' 2026 rapid review of twenty-nine ambulatory studies found moderate accuracy for total sleep time and time in bed, with weaker estimates for sleep efficiency, wake after sleep onset and stages; REM and deep-sleep estimates were especially unreliable, and clinical populations were rarely included. Khosla and colleagues' AASM statement remains the clinical boundary: consumer technologies should not diagnose or treat sleep disorders without appropriate validation and regulatory status. Baron and colleagues introduced *orthosomnia* through clinical cases; it is descriptive rather than a formal diagnosis or prevalence estimate.

Current verification. Current guideline, institutional and research status was checked on 2 September 2026. Changeable claims are labelled by the evidence and date available rather than presented as permanent settlement.

Bibliography

Guidelines and official references

American Academy of Sleep Medicine and Sleep Research Society.

“Recommended Amount of Sleep for a Healthy Adult: A Joint Consensus Recommendation.” *Journal of Clinical Sleep Medicine* 11, no. 6 (2015): 591-592. DOI: 10.5664/jcsm.4758.

Auger, Robert R., Helen J. Burgess, Jonathan S. Emens, Leslie V. Deriy, Sherene M. Thomas, and Kimberly M. Sharkey. “Clinical Practice Guideline for the Treatment of Intrinsic Circadian Rhythm Sleep-Wake Disorders: Advanced Sleep-Wake Phase Disorder, Delayed Sleep-Wake Phase Disorder, Non-24-Hour Sleep-Wake Rhythm Disorder, and Irregular Sleep-Wake Rhythm Disorder. An Update for 2015.” *Journal of Clinical Sleep Medicine* 11, no. 10 (2015): 1199-1236. DOI: 10.5664/jcsm.5100.

Buyse, Daniel J., J. Todd Arnedt, Luis Buenaver, et al. “Combination

Treatment for Chronic Insomnia Disorder in Adults: An American Academy of Sleep Medicine Clinical Practice Guideline.” *Journal of Clinical Sleep Medicine* 22 (2026): article 56. Published 13 April 2026. DOI: 10.1007/s44470-025-00038-8.

Edinger, Jack D., J. Todd Arnedt, Suzanne M. Bertisch, et al. “Behavioral and Psychological Treatments for Chronic Insomnia Disorder in Adults: An American Academy of Sleep Medicine Clinical Practice Guideline.” *Journal of Clinical Sleep Medicine* 17, no. 2 (2021): 255-262. DOI: 10.5664/jcsm.8986.

National Institute of Neurological Disorders and Stroke. *Narcolepsy*. Bethesda, MD: National Institutes of Health, accessed 2 September 2026.

National Heart, Lung, and Blood Institute. *Sleep Apnea*. Bethesda, MD: National Institutes of Health, updated 9 January 2025.

National Heart, Lung, and Blood Institute. *Sleep Deprivation and Deficiency*. Bethesda, MD: National Institutes of Health, updated 24 March 2022.

National Institute of General Medical Sciences. *Circadian Rhythms*. Bethesda, MD: National Institutes of Health, updated 20 May 2025.

National Institute for Health and Care Excellence. *Menopause: Identification and Management*. NICE guideline NG23. London: NICE, updated 15 April 2026; recommendations checked 2 September 2026.

National Institute for Health and Care Excellence. *Obstructive Sleep Apnoea/Hypopnoea Syndrome and Obesity Hypoventilation Syndrome in Over 16s*. NICE guideline NG202. London: NICE, 2021.

Riemann, Dieter, Colin A. Espie, Ellemarije Altena, et al. “The European Insomnia Guideline: An Update on the Diagnosis and Treatment of Insomnia 2023.” *Journal of Sleep Research* 32, no. 6 (2023): e14035. DOI: 10.1111/jsr.14035.

St-Onge, Marie-Pierre, Brooke Aggarwal, Julio Fernandez-Mendoza, et al. “Multidimensional Sleep Health: Definitions and Implications for

Cardiometabolic Health: A Scientific Statement From the American Heart Association.” *Circulation: Cardiovascular Quality and Outcomes* 18, no. 5 (2025): e000139. DOI: 10.1161/HCQ.0000000000000139.

Research and reviews

Aserinsky, Eugene, and Nathaniel Kleitman. “Regularly Occurring Periods of Eye Motility, and Concomitant Phenomena, during Sleep.” *Science* 118, no. 3062 (1953): 273-274. DOI: 10.1126/science.118.3062.273.

Baron, Kelly Glazer, Sabra Abbott, Nancy Jao, Natalie Manalo, and Rebecca Mullen. “Orthosomnia: Are Some Patients Taking the Quantified Self Too Far?” *Journal of Clinical Sleep Medicine* 13, no. 2 (2017): 351-354. DOI: 10.5664/jcsm.6472.

Besedovsky, Luciana, Tanja Lange, and Monika Haack. “The Sleep-Immune Crosstalk in Health and Disease.” *Physiological Reviews* 99, no. 3 (2019): 1325-1380. DOI: 10.1152/physrev.00010.2018.

Birrer, Vera, Mohamed Elgendi, Olivier Lambercy, et al. “Evaluating Reliability in Wearable Devices for Sleep Staging.” *npj Digital Medicine* 7 (2024): 74. DOI: 10.1038/s41746-024-01016-9.

Borbély, Alexander A., Serge Daan, Anna Wirz-Justice, and Tom Deboer. “The Two-Process Model of Sleep Regulation: A Reappraisal.” *Journal of Sleep Research* 25, no. 2 (2016): 131-143. DOI: 10.1111/jsr.12371.

Carskadon, Mary A. “Sleep in Adolescents: The Perfect Storm.” *Pediatric Clinics of North America* 58, no. 3 (2011): 637-647. DOI: 10.1016/j.pcl.2011.03.003.

Chang, Anne-Marie, Daniel Aeschbach, Jeanne F. Duffy, and Charles A. Czeisler. “Evening Use of Light-Emitting eReaders Negatively Affects Sleep, Circadian Timing, and Next-Morning Alertness.” *Proceedings of the National Academy of Sciences* 112, no. 4 (2015): 1232-1237. DOI: 10.1073/pnas.1418490112.

Depner, Christopher M., Edward L. Melanson, Robert H. Eckel, et al. “Ad

Libitum Weekend Recovery Sleep Fails to Prevent Metabolic Dysregulation during a Repeating Pattern of Insufficient Sleep and Weekend Recovery Sleep.” *Current Biology* 29, no. 6 (2019): 957-967.e4. DOI: 10.1016/j.cub.2019.01.069.

Dagum, Paul, Donald L. Elbert, Laurent Giovangrandi, et al. “The Glymphatic System Clears Amyloid Beta and Tau from Brain to Plasma in Humans.” *Nature Communications* 17 (2026): 715. DOI: 10.1038/s41467-026-68374-8.

Gardiner, Carissa L., Jonathon Weakley, Louise M. Burke, et al. “Dose and Timing Effects of Caffeine on Subsequent Sleep: A Randomized Clinical Crossover Trial.” *Sleep* 48, no. 4 (2025): zsae230. DOI: 10.1093/sleep/zsae230.

Gardiner, Carissa, Jonathon Weakley, Louise M. Burke, Gregory D. Roach, Charli Sargent, Nirav Maniar, Minh Huynh, Dean J. Miller, Andrew Townshend, and Shona L. Halson. “The Effect of Alcohol on Subsequent Sleep in Healthy Adults: A Systematic Review and Meta-Analysis.” *Sleep Medicine Reviews* 80 (2025): 102030. DOI: 10.1016/j.smr.2024.102030.

Haghighy, Shahab, Sepideh Khoshnevis, Michael H. Smolensky, Kenneth R. Diller, and Richard J. Castriotta. “Before-Bedtime Passive Body Heating by Warm Shower or Bath to Improve Sleep: A Systematic Review and Meta-Analysis.” *Sleep Medicine Reviews* 46 (2019): 124-135. DOI: 10.1016/j.smr.2019.04.008.

Hauglund, Natalie L., Mie Andersen, Klaudia Tokarska, et al. “Norepinephrine-Mediated Slow Vasomotion Drives Glymphatic Clearance during Sleep.” *Cell* 188, no. 3 (2025): 606-622.e17. DOI: 10.1016/j.cell.2024.11.027.

He, Ying, Christopher R. Jones, Nobuhiro Fujiki, et al. “The Transcriptional Repressor DEC2 Regulates Sleep Length in Mammals.” *Science* 325, no. 5942 (2009): 866-870. DOI: 10.1126/science.1174443.

Jones, Samuel E., Jacqueline M. Lane, Andrew R. Wood, et al.

“Genome-Wide Association Analyses of Chronotype in 697,828 Individuals Provides Insights into Circadian Rhythms.” *Nature Communications* 10 (2019): 343. DOI: 10.1038/s41467-018-08259-7.

Khosla, Seema, Maryann C. Deak, Dennis Gault, et al. “Consumer Sleep Technology: An American Academy of Sleep Medicine Position Statement.” *Journal of Clinical Sleep Medicine* 14, no. 5 (2018): 877-880. DOI: 10.5664/jcsm.7128.

Landvatter, Joshua, Jeanna T. Ryan, Christopher M. Depner, and Kelly Baron. “Real-World Use of Consumer Sleep Devices: A Rapid Review.” *Chest* 169, no. 4 (2026): 1101-1110. DOI: 10.1016/j.chest.2025.10.039.

Loomis, Alfred L., E. Newton Harvey, and Garret A. Hobart. “Cerebral States during Sleep, as Studied by Human Brain Potentials.” *Journal of Experimental Psychology* 21, no. 2 (1937): 127-144. DOI: 10.1037/h0057431.

Miao, Andawei, Tianyuan Luo, Bryan Hsieh, et al. “Brain Clearance Is Reduced during Sleep and Anesthesia.” *Nature Neuroscience* 27, no. 6 (2024): 1046-1050. DOI: 10.1038/s41593-024-01638-y.

Rasch, Björn, and Jan Born. “About Sleep's Role in Memory.” *Physiological Reviews* 93, no. 2 (2013): 681-766. DOI: 10.1152/physrev.00032.2012.

Reichert, Carolin F., Tom Deboer, and Hans-Peter Landolt. “Adenosine, Caffeine, and Sleep-Wake Regulation: State of the Science and Perspectives.” *Journal of Sleep Research* 31, no. 4 (2022): e13597. DOI: 10.1111/jsr.13597.

Schyvens, An-Marie, Brent Peters, Nina Catharina Van Oost, et al. “A Performance Validation of Six Commercial Wrist-Worn Wearable Sleep-Tracking Devices for Sleep Stage Scoring Compared to Polysomnography.” *SLEEP Advances* 6, no. 2 (2025): zpaf021. DOI: 10.1093/sleepadvances/zpaf021.

Skorucak, Jelena, Emma L. Arbon, Derk-Jan Dijk, and Peter Achermann. “Response to Chronic Sleep Restriction, Extension, and Subsequent Total

Sleep Deprivation in Humans: Adaptation or Preserved Sleep Homeostasis?" *Sleep* 41, no. 7 (2018): zsy078. DOI: 10.1093/sleep/zsy078.

Sletten, Tracey L., et al. "The Importance of Sleep Regularity: A Consensus Statement of the National Sleep Foundation Sleep Timing and Variability Panel." *Sleep Health* 9, no. 6 (2023): 801-820. DOI: 10.1016/j.sleh.2023.07.016.

Stutz, Jan, Remo Eiholzer, and Christina M. Spengler. "Effects of Evening Exercise on Sleep in Healthy Participants: A Systematic Review and Meta-Analysis." *Sports Medicine* 49, no. 2 (2019): 269-287. DOI: 10.1007/s40279-018-1015-0.

Van Dongen, Hans P. A., Greg Maislin, Janet M. Mullington, and David F. Dinges. "The Cumulative Cost of Additional Wakefulness: Dose-Response Effects on Neurobehavioral Functions and Sleep Physiology from Chronic Sleep Restriction and Total Sleep Deprivation." *Sleep* 26, no. 2 (2003): 117-126. DOI: 10.1093/sleep/26.2.117.

Winkelman, John W., Joshua A. Berkowski, Lourdes M. DelRosso, et al. "Treatment of Restless Legs Syndrome and Periodic Limb Movement Disorder: An American Academy of Sleep Medicine Clinical Practice Guideline." *Journal of Clinical Sleep Medicine* 21, no. 1 (2025): 137-152. DOI: 10.5664/jcsm.11390.

Wittmann, Marc, Jenny Dinich, Martha Merrow, and Till Roenneberg. "Social Jetlag: Misalignment of Biological and Social Time." *Chronobiology International* 23, nos. 1-2 (2006): 497-509. DOI: 10.1080/07420520500545979.

Xie, Lulu, Hongyi Kang, Qiwu Xu, et al. "Sleep Drives Metabolite Clearance from the Adult Brain." *Science* 342, no. 6156 (2013): 373-377. DOI: 10.1126/science.1241224.

Books

Espie, Colin. *Overcoming Insomnia 2nd Edition*. London: Robinson, 2021.

Kryger, Meir. *The Mystery of Sleep: Why a Good Night's Rest Is Vital to a Better, Healthier Life*. New Haven: Yale University Press, 2017.

Leschziner, Guy. *The Secret World of Sleep: Journeys Through the Nocturnal Mind*. London: Simon & Schuster UK, 2020.

Roenneberg, Till. *Internal Time: Chronotypes, Social Jet Lag, and Why You're So Tired*. Cambridge, MA: Harvard University Press, 2012.