

THE IN A HURRY SERIES

The Ice Age

in a Hurry

When the world froze, and what survived

Max Brocklesby

BOOKSINAHURRY.COM

Contents

The Whole Thing in One Page 4

Why You Should Care 5

The Core Ideas 7

How It Actually Works 20

What People Get Wrong 29

Use It 34

Terms 39

Go Deeper 43

Notes and Sources 45

Bibliography 51

The Whole Thing in One Page

The Ice Age is usually pictured as one long white emergency. Mammoths cross a blizzard, people crouch in caves, a wall of ice reaches south, and then the world warms and resumes normal service. Almost every part of that picture is misleading.

There was no single freeze. During the Pleistocene, from 2.58 million to 11,700 years ago, great ice sheets repeatedly advanced and retreated. Cold glacials alternated with warmer interglacials. Even at the Last Glacial Maximum, roughly 26,500 to 19,000 or 20,000 years ago, most of Earth was ice-free. The global mean was about 6 degrees Celsius colder than the pre-industrial world, but the difference was distributed unevenly. Northern continents changed far more than many tropical regions, and cold often arrived with dryness.

Small shifts in Earth's orbit paced the cycles by changing where and when sunlight arrived. The decisive season was often summer. Snow that survives a cool summer can accumulate into ice; snow that melts cannot, however brutal the preceding winter. Ice then reflected more sunlight. Carbon dioxide fell, oceans and winds changed, dust moved, and the climate response grew far larger than the original orbital nudge. Orbit kept time. The Earth system supplied the volume.

The resulting ice did more than lower temperature. It stored ocean water on land, pushing global sea level roughly 125 to 130 metres below today's near its minimum. Britain joined continental Europe across Doggerland. Siberia and Alaska met across Beringia. Rivers were blocked or redirected, crust bent beneath the weight, and coastlines moved hundreds of kilometres in some places. Ice was a geography machine.

Life did not march south as one obedient biome. The mammoth steppe was a varied belt of cold, dry grassland, herbs and shrubs, with no exact modern equivalent. Woolly mammoths, horses, bison, reindeer, saiga and predators shared different parts of it. Forests contracted into refugia. Cold species later did the same when warmth returned. Each population met a different

map, and survival depended on whether suitable habitat remained connected, whether a refuge held, how quickly animals bred, and whether they could move or alter their behaviour.

Humans were inside this system, not watching from its edge. Neanderthals, Denisovans and Homo sapiens endured repeated climatic swings through fire, tools, clothing, shelters, movement, social learning and cooperation. They also met and interbred. A vanished population can therefore survive partly in another population's genome.

The final thaw was no gentle release. Warming came in jumps, ice sheets collapsed over millennia, seas rose, habitats fragmented, and the North Atlantic briefly returned to severe cold during the Younger Dryas. Large animals disappeared at different times on different continents. Expanding human pressure mattered strongly, sometimes decisively, while habitat change, small populations and slow reproduction altered the outcome. There is no single global verdict.

What survived was selected by routes, refuges, timing and chance. What remains is under your feet, in the shape of lakes and valleys, in living genomes, and in the permanent polar ice that means the wider icehouse has not entirely gone away.

That is the book.

Why You Should Care

Stand on the east coast of Britain and look towards the North Sea. During the coldest part of the last glacial cycle, much of the water in front of you was not there. It had been lifted out of the ocean, carried as vapour and snow, and stored in ice sheets. Across the long interval when the shelf stood exposed, its plains held rivers, marshes and grazing animals and, when conditions allowed, people. Britain was the western edge of a continent, not an island. Fishing and aggregate work have recovered animal bones and worked stone from the seabed, physical reminders that a modern shipping lane can occupy an old human landscape.

That vanished country is called Doggerland. Its loss gives the Ice Age its first claim on your attention: climate does not merely alter the weather over a fixed map. It can move the map itself. Coasts advance and drown. Rivers find new courses. Mountain ice grinds rock into sediment, then leaves lakes, ridges and unstable slopes behind. Land depressed by an ice sheet continues to rise after the weight has gone. Much of the physical world that feels permanent is the residue of water changing address.

The second reason is biological. The animals and plants around you are not a balanced local assembly that has occupied the same places since deep time. They are survivors of repeated separation, retreat and return. A tree species may carry genetic divisions formed in different glacial refugia. Two populations may meet along an old expansion front. A cold-adapted animal can become stranded on a mountain as the lowlands warm, while a temperate species recolonises ground that ice had erased. The present distribution is a frame from a long film of movement.

This changes the question you ask about survival. The obvious question is how a mammoth kept warm. The more useful one is whether its food remained reachable, whether winter snow became too deep, whether populations stayed connected, and whether enough calves survived to replace adults. Fur solves one problem. It cannot reopen a lost corridor or make a slow-breeding population large again.

Humans belong in the same account. The Ice Age was not scenery behind human evolution. It repeatedly divided populations, opened routes, closed them, concentrated people in workable regions and placed different human groups in contact. Neanderthals and modern humans did not occupy separate museum cases. They met, competed, interbred and left descendants whose genomes preserve part of that encounter. Our species survived through bodies, technology and collective knowledge, then became one of the pressures other species had to survive.

The period also offers a clean lesson in how large systems change. Earth's orbital variations are modest and predictable. The resulting climate swings were neither modest nor mechanically proportional because ice, vegetation,

greenhouse gases and oceans fed back on the initial change. A pacemaker sets the timing; feedback determines the scale. Confusing trigger with amplifier is one of the quickest ways to misunderstand any complex system.

There are limits to the comparison with the present. The glacial cycles unfolded under different starting conditions, over far longer spans, and were paced by orbital changes rather than modern industrial emissions. The Ice Age is evidence that climate can change, not an excuse to treat every climate change as the same event.

Its deeper value is less argumentative. It lets you see familiar ground as temporary, living communities as historical, and survival as a problem of timing and connection. The blizzard is memorable. The moving coastline, shrinking refuge and broken population are what decided the outcome. Once those are visible, the frozen world becomes stranger, more inhabited and far more useful to understand.

The Core Ideas

The Ice Age Was a Pulse, Not a Pause

The phrase Ice Age encourages the wrong shape. It sounds like an event with a beginning, a frozen middle and an ending. The Pleistocene was closer to a pulse: repeated expansions and contractions of land ice, nested within a much longer cool state of the planet. Large Antarctic ice sheets began forming about 34 million years ago, and major Northern Hemisphere glaciation intensified around 2.7 million years ago. The Pleistocene is the most recent and humanly familiar chapter, not the whole history of planetary ice.

Geologists place the start of the Pleistocene at 2.58 million years ago and its end at 11,700 years ago. That interval contains many glacials, when ice sheets expanded and global sea level fell, and interglacials, when much of the ice retreated and seas rose. We live in one of the warmer intervals, the Holocene. The popular singular, the Ice Age, is therefore useful only if it is understood as a family of cycles rather than one continuous freeze.

Even the cold phases were not uniformly white. At the Last Glacial Maximum, nearly all major ice sheets stood near their greatest global extents from about 26,500 to 19,000 or 20,000 years ago. The Laurentide ice sheet covered much of Canada and reached into the northern United States. The Cordilleran ice sheet occupied western mountains. A Fennoscandian sheet spread across northern Europe, while mountain glaciers enlarged from the Alps to New Zealand and Patagonia. Yet Africa, most of Asia, Australia and wide tropical belts remained free of continental ice.

The global mean temperature was about 6 degrees Celsius below the pre-industrial mean. That number is large for a planetary average and small compared with some local changes. High northern latitudes cooled far more. Tropical oceans cooled less. Many regions became drier because colder air carried less moisture and altered circulation reduced rainfall. Some places that escaped ice still became treeless, dusty or seasonally severe. The Ice Age was a reorganisation of heat and water, not a coat laid evenly across a globe.

The cycles also changed through time. For much of the early Pleistocene, glacial variation followed a strong rhythm near 41,000 years, matching changes in Earth's axial tilt. Between roughly 1.25 million and 750,000 years ago, the system shifted towards larger, longer cycles averaging close to 100,000 years. No new 100,000-year orbital force appeared with enough strength to explain the change. Ice sheets, bed conditions, oceans and the carbon cycle began responding differently. This Mid-Pleistocene Transition remains one of the central unsolved problems in palaeoclimate.

That uncertainty matters because it blocks a clockwork picture. The orbit did not dictate a fixed amount of ice. It altered the conditions under which a responsive Earth system grew or lost ice. Previous ice sheets changed soils and rock beneath them. Carbon moved among ocean, atmosphere and land. The size of one cycle affected the starting point of the next.

The first mental correction is therefore temporal. Do not picture a frozen world waiting millions of years for release. Picture a world that repeatedly

moved between states, never returning to precisely the same geography or ecology. Survival meant enduring not one cold extreme but the movement between cold, less cold, dry, wet, connected and fragmented conditions.

Summer Sunlight Set the Pace

A spectacular winter can kill animals, close roads and cover a continent in snow. It cannot build an ice sheet by itself. Continental ice grows when more snow survives one year than is lost before the next accumulation season. That makes summer melt as important as winter snowfall, and often more decisive.

Earth's path around the Sun changes in three main ways. The orbit becomes slightly more or less elliptical, called eccentricity. The tilt of Earth's axis varies, called obliquity. The axis also wobbles, called precession, changing which season falls at which part of the orbit. None of these variations changes the Sun's total output. They redistribute sunlight by season and latitude. Tilt varies on a rhythm near 41,000 years, while precession produces shorter rhythms near 19,000 to 23,000 years. Eccentricity alters how strongly precession matters and carries a prominent rhythm near 100,000 years, but its direct global forcing is weak. That mismatch is part of the famous 100,000-year problem.

Northern summer sunlight has unusual leverage because the Northern Hemisphere holds large continents at high and middle latitudes. If summer sunlight weakens, winter snow is more likely to persist across land. Surviving snow reflects much of the next summer's energy. More snow can then survive, compact into firn, become ice and begin to flow under its own weight. If summer sunlight strengthens, melting attacks the same system. The calendar of sunlight sets conditions for growth or retreat.

The orbital changes are too weak to explain glacial scale alone. Their power lies in feedback. Bright snow and ice replace darker land or ocean and reflect more solar energy, adding cooling. Vegetation shifts, changing reflectivity, dust and carbon storage. Cooler oceans can hold more dissolved carbon dioxide, while circulation and biological activity change

how carbon moves. Atmospheric carbon dioxide and methane fell during glacials, reducing greenhouse warming. Expanding ice sheets also rose into colder air and altered winds. Each response changed the next response.

The sequence is not a neat chain in which orbit acts, carbon follows and ice obeys. Timing varies among hemispheres and variables. Antarctic temperature, greenhouse gases, ocean circulation and northern ice are coupled parts of one system. Ice-core records show carbon dioxide moving closely with Antarctic climate over repeated cycles, while marine sediments record the changing volume of global ice. Carbon dioxide often amplified changes that orbital geometry helped initiate, then became a forcing in its own right as its concentration changed.

This distinction between pacemaker and amplifier prevents two opposite errors. One is to say orbital cycles caused everything, as though kilometres of ice were a direct shadow cast by a small change in sunlight. The other is to say greenhouse gases cannot matter because temperature sometimes began changing before carbon dioxide. A feedback can respond to an initial change and then enlarge it. Cause is a role in a sequence, not a title awarded to the first variable that moved.

There were also thresholds and delays. An ice sheet can persist through several unfavourable summers because of its size and height. Meltwater can alter ocean circulation. Bedrock and sediment beneath the ice can change how fast it flows. The carbon cycle responds on several timescales. Glacial terminations were therefore much faster than the long build-up that preceded them, while individual regions could jump abruptly even during a slow orbital trend.

The useful picture is a small seasonal shove applied to a machine full of memory. The orbit supplied regular changes in timing. Feedbacks, thresholds and inherited conditions decided how much world was remade.

Ice Moved the Map

An ice sheet is stored ocean. Snow falls on land, survives, compacts and remains there for centuries or millennia. When continental ice grows, the sea must fall. Sea ice is different: it already floats, so freezing more of it has little direct effect on global sea level. Near the Last Glacial Maximum, global sea level reached roughly 125 to 130 metres below today's level. That change exposed continental shelves, joined islands to mainlands and shifted some coasts far beyond the modern horizon.

Beringia linked northeast Asia and northwest North America across a broad, cold region. It was not a narrow causeway used once by travellers hurrying between continents. For long periods it was a landscape with rivers, vegetation, animals and human populations, while ice and climate controlled which routes opened beyond it. Doggerland joined Britain to continental Europe across what is now the southern North Sea. Southeast Asia's exposed Sunda Shelf connected many present islands. Australia and New Guinea formed the larger landmass of Sahul, though deep-water channels still separated it from Asia.

These connections rearranged evolution and human movement without guaranteeing either. A land bridge is useful only if it carries suitable habitat and leads somewhere passable. Beringia could connect steppe populations while continental ice blocked routes farther south. Falling seas might unite populations in one place and isolate marine populations elsewhere. When water returned, the same lowlands became traps, estuaries and seabed.

The weight of ice changed solid ground. Kilometres of ice depressed the crust and pushed the mantle beneath it aside. Once the load melted, land began to rebound, a process still continuing in formerly glaciated regions. Areas around the former ice margins can sink as mantle material returns towards the rising centre. Relative sea level therefore varies by place: the ocean rises globally while land itself moves up or down.

Ice also redirected water before it became sea. A sheet can dam rivers, force drainage along its margin and create immense lakes. Outburst floods

cut channels and moved sediment in hours or days. Repeated glaciation deepened some basins, scoured rock, carried boulders far from their source and spread till, sand and gravel. The Great Lakes occupy basins shaped by ice acting on older geology. Scandinavia, Canada, Scotland and the Alps carry the same signature in different forms: polished rock, U-shaped valleys, moraines, drumlins, erratics and chains of lakes.

Beyond the ice margin, wind did another kind of earth-moving. Sparse vegetation and glacial grinding supplied fine sediment. Winds carried this dust and deposited thick blankets of loess across parts of Europe, Asia and North America. Those deposits later weathered into productive soils in many regions. Ice-age cold could therefore help create future farmland far from the ice itself, though erosion, mineral composition and later climate determined the result.

The map changed above ground as well. Vast ice sheets altered atmospheric circulation. Colder oceans, expanded sea ice and exposed shelves shifted storm tracks and rainfall. Deserts expanded in some regions. Tropical mountain glaciers descended. Monsoon systems weakened or moved. The label glacial can hide opposite local experiences: buried beneath ice, exposed as dry steppe, cooled rainforest, enlarged desert, wetter lake basin or windy tundra.

A modern political map makes these changes hard to imagine because it treats coastlines as borders of reality. During the Pleistocene they were moving contours in a water budget. The ice sheets did not occupy empty northern margins. They repeatedly rebuilt the routes by which species met, separated and survived.

Cold Built Ecosystems With No Modern Address

The woolly mammoth is usually placed in a snowy wasteland, as though large herbivores fed on whiteness. Its main Eurasian and Beringian habitat was the mammoth steppe, a broad but varied mosaic of grasses, sedges, flowering herbs and low shrubs under cold, dry conditions. It stretched across regions that are now divided among tundra, boreal forest and temperate landscapes. No exact modern biome reproduces it.

Its fame is also partly an archive effect. Large bones survive, tusks are conspicuous, and frozen ground can preserve hair, skin and stomach contents that would disappear elsewhere. The Ice Age bestiary therefore looks richer in detail than many warmer ecosystems, although preservation remains patchy and biased towards particular places and bodies.

Dryness was central. Heavy, persistent snow can seal forage away from grazing animals. Cold steppe with shallow snow may be easier to feed in than a milder winter under deep wet cover. Low precipitation, wind and grazing helped keep some ground open. In summer, long daylight and nutrient-rich soils could support abundant plant growth. Permafrost restricted drainage and roots in some places, while windblown loess supplied fresh mineral material in others. Mammoths, horses, bison, reindeer, saiga, woolly rhinoceroses and musk oxen occupied different portions and periods of this system, accompanied by predators and scavengers. They never formed one fixed cast across an unchanging stage.

Ancient environmental DNA has sharpened the picture. Sediment from Arctic lakes and permafrost contains fragments shed by plants and animals. A large circumpolar study of samples spanning the past 50,000 years found regionally varied steppe-tundra communities during the Last Glacial Maximum, followed by increasing divergence among regions as climate warmed. Grasses alone did not dominate everywhere. Forbs, willows and other plants mattered, and wetter conditions were associated with declining diversity of large grazing animals in the studied Arctic regions.

That evidence corrects the old image while carrying its own boundary. Arctic

sediment sites do not settle the cause of every extinction on every continent. Environmental DNA can show that a taxon was present near a sampled place, but abundance, precise range and ecological importance require other evidence. The mammoth steppe remains a useful name for a family of cold open ecosystems, not a single carpet rolled across the north.

Large herbivores may also have helped maintain open conditions by grazing, trampling, moving nutrients and damaging young trees. The effect is plausible and supported in some settings, but climate and moisture set strong limits. Mammoths did not create their own continent by force of appetite. Ecosystems arise from interaction among climate, soils, plants, herbivores, predators, fire and history, with the balance differing across space.

When warmth returned, the old community did not move north as a unit. Trees expanded at different rates. Wetlands spread. Some grazing species shifted, some contracted into small ranges, some changed diet, and some disappeared. Reindeer and musk ox survived. Woolly rhinoceroses did not. Mammoth populations lingered in separated pockets long after their mainland range collapsed.

A biome name is therefore a convenient summary, not a travelling organism. Climate change dissolved old combinations and assembled new ones from unequal survivors. What vanished was more than a list of species. It was a set of relationships that the modern world no longer places together.

Survival Happened in Corridors and Refugia

Cold tolerance is visible, so it attracts explanation. Thick fur, fat, compact bodies and seasonal coats matter. Yet a population can possess all of them and still disappear if its habitat becomes too small, too isolated or too unreliable. Survival is a demographic achievement performed on a map.

A refugium is a place where a population persists while conditions become unsuitable across much of its former range. Temperate trees survived glacial periods in workable pockets, often in southern Europe but not only there. Local topography, coastal influence, groundwater, slope, fire and

microclimate could preserve habitat farther north than a simple line on a map suggests. Cold-adapted species faced the reverse problem during interglacials, retreating towards high latitudes, mountains or continental interiors.

There is no universal refuge. One species needs woodland, another open ground, another a host plant, another snow of a particular depth. A valley that protects trees from cold may trap a grazing animal behind forest. A mountain can preserve cool climate while leaving too little area for a large population. Refugia vary in size and duration, and the population inside them may lose genetic diversity even when the species survives. They are also temporary by definition. A refuge from glacial cold may become unsuitable during the following warm interval, so persistence can require repeated shifts among different safe areas rather than loyalty to one sanctuary.

Corridors determine what happens next. When suitable conditions expand, populations can recolonise. Fast-moving animals may track habitat across generations. Trees move through seed dispersal and recruitment, which can be rapid over centuries but slow beside abrupt climate shifts. Rivers, mountain chains, ice margins, coastlines and exposed shelves can guide movement or stop it. A route that connects two populations may permit gene flow; flooding the same route can isolate them.

These repeated contractions and expansions left genetic structure in living species. Populations separated in different refugia accumulate different variants through mutation, selection and drift. When they expand and meet, their descendants can form contact zones or mix. Some lineages vanish during the contraction, leaving a narrower sample to repopulate a large area. The modern range may look broad while carrying the signature of a past bottleneck.

Fossils and ancient DNA show that responses were individualistic. Woolly mammoth, woolly rhinoceros, horse, reindeer, bison and musk ox did not rise and fall together under one climate switch. Habitat suitability, human presence and population size affected each differently. Even members of

one species could have distinct regional histories. A continental label such as glacial cooling is too coarse to predict a local fate.

Mobility itself has limits. Following climate requires suitable ground to move into, time to reproduce there and enough connected individuals to avoid demographic collapse. A range can shift only as fast as the organism and its ecological partners permit. A predator may depend on prey that moves differently. A plant may outrun one pollinator and lose another. Disease and competitors can arrive through the same corridor that offers escape.

This is why survival should be separated into levels. A species can persist while many populations disappear. A lineage can survive through admixture while its distinct population ends. A forest type can return without the same species mixture. A landscape can remain green while its Ice Age genetic diversity has been reduced.

The lesson is not that refugia guarantee safety. They buy continuity. During repeated glacial pulses, continuity was enough to turn a small hidden population into the source of a later continental return. Where no refuge remained, adaptation had no population left on which to work.

Humans Were Ice Age Animals

The familiar Ice Age human is a modern person in fur, inserted into a hostile landscape as the clever exception to nature. The better starting point is that several human populations were native components of Pleistocene ecosystems. They hunted, gathered, scavenged, moved, raised children, cared for injured people and altered local environments under the same constraints as other large mammals.

Neanderthals occupied western Eurasia through repeated climatic swings. Denisovans are known most clearly from DNA and sparse remains across Asia. *Homo sapiens* evolved in Africa and expanded through regions that ranged from tropical coast to cold steppe. These groups were not sealed species in separate chapters. Genomes show that they met and interbred. Many living people outside Africa retain Neanderthal ancestry, while Denisovan ancestry is especially substantial in some populations of

Oceania and parts of Asia. A population can vanish as a distinct group while part of it continues biologically elsewhere.

Human survival did not rest on one invention. Fire supplied heat, light, food processing and protection. Stone, bone and antler tools extended hands and teeth. Shelters reduced exposure. Clothing increased the climates a body could occupy, though most early garments have decayed and their history must be reconstructed indirectly. Hunting and gathering knowledge turned seasonal movement into a plan rather than a flight. Social learning allowed a technique discovered by one person to outlive that person.

The decisive adaptation was collective and cumulative. A lone human is a poor Ice Age animal. A group can divide tasks, share food, care for children, remember distant water and exchange materials or partners with other groups. Wider social networks can buffer a local failure: information, stone, ornaments and genes move beyond one camp. Burials, pigments, carvings and cave art show that some communities also sustained symbolic worlds under severe conditions. Those practices were not decorative proof of intelligence; they belonged to relationships through which identity, memory and obligation travelled. Archaeological distributions show long-distance connections in some Upper Palaeolithic societies, but network size and resilience varied greatly through time and place.

This does not produce a clean contest in which modern humans were smart and Neanderthals were not. Neanderthals used fire, made complex tools, hunted demanding prey and cared for group members. Their disappearance by around 40,000 years ago was regionally staggered. Small and fragmented populations may have been vulnerable to climatic instability, competition, disease, low fertility or chance. Some were absorbed through interbreeding. The evidence does not isolate one cause with the confidence required for a superiority fable.

Homo sapiens also failed often. Early expansions left little or no ancestry in later populations. Communities disappeared when routes closed or resources changed. Survival of the species conceals repeated local endings. The groups that endured were not representatives of an inevitable

winner. They were populations whose bodies, knowledge, alliances and timing happened to work under particular conditions.

By the late Pleistocene, humans were also changing the survival problem for other animals. Hunting can have large effects on slow-breeding species even when kills are infrequent. Fire, disturbance and competition may alter habitat. Human pressure did not have the same intensity everywhere, and climate change could reduce populations before people delivered an additional loss. Yet humans cannot be treated as another neutral climate proxy. They learned, cooperated and targeted prey, so their impact could grow while their own numbers remained modest.

The Ice Age made human culture matter because conditions moved faster than genetic adaptation alone could answer. Culture made humans resilient. It also made them an unusually mobile source of risk.

The Thaw Was Another Extinction Test

The end of a glacial period sounds like rescue. Temperatures rise, ice retreats and land becomes habitable again. For organisms fitted to cold, dry, open country, the same sequence can remove the world they require.

Northern deglaciation began around 19,000 to 20,000 years ago as summer sunlight increased. Ice loss then interacted with greenhouse gases, oceans and changing reflectivity. Global sea level rose by more than a hundred metres over the long transition. Lowlands flooded. Ice-margin lakes formed and drained. Forests expanded into former steppe. Rainfall and snow depth changed. Populations that had survived the cold were divided among islands, mountains and shrinking continental pockets.

The warming was uneven and interrupted. During the Bølling-Allerød interval, North Atlantic regions warmed sharply. Around 12,900 years ago, the Younger Dryas brought a severe return of cold conditions to the North Atlantic region, lasting until about 11,700 years ago. It was not a second global Last Glacial Maximum. Southern extratropical temperatures did not mirror the northern event, and tropical responses varied. Changes in Atlantic overturning circulation, atmospheric circulation and radiative forcing

interacted; no single mechanism reproduces the whole pattern.

This instability met increasingly widespread human populations. Late Quaternary losses removed many large mammals, but not in one simultaneous wave. Australia experienced major losses earlier than the Americas. Eurasian extinctions stretched across a long interval. North and South America lost many large-bodied genera near the end of the Pleistocene, after people were present. Islands later produced their own human-linked extinctions under different conditions.

Evidence gives humans a major role without granting one universal mechanism. The timing of human arrival and extinction severity are strongly associated at broad scales. Archaeological kill sites prove hunting of some species, not every species. Slow reproduction means even limited additional mortality can push a small population below recovery. Climate and habitat change also altered ranges, productivity and connectivity. Their effects were species-specific, and a population already compressed by habitat loss could become far more sensitive to hunting.

The woolly mammoth shows why a clean date misleads. Mainland populations contracted as open cold habitat fragmented, yet isolated populations survived for millennia. Mammoths persisted on Wrangel Island until roughly 4,000 years ago, long after the Pleistocene had ended. Their final disappearance cannot explain the earlier continental collapse, and the continental collapse cannot be reduced to the last island's fate. Extinction is a process distributed across populations.

Size mattered, but not alone. Large animals tend to reproduce slowly, need large ranges and exist at lower densities. Specialised diets, migration routes and dependence on open habitat could add risk. Smaller animals disappeared too, while some giants survived. Reindeer, bison, musk oxen and large predators reached the modern world in reduced or shifted ranges. Outcome depended on ecological flexibility, geography, human pressure and chance.

The Holocene boundary at 11,700 years ago marks the end of the

Pleistocene, not a return to a pre-Ice Age Earth. Coastlines continued moving, ice-loaded land continued rebounding, soils developed on glacial sediment and human populations expanded through inherited landscapes. Antarctica and Greenland retained permanent ice. In the broad geological sense, Earth remains in an icehouse state.

The final thaw shows why repeated movement was the defining pressure. Warmth was another movement, not the cancellation of change. The Ice Age filtered survivors twice: through the cold, then through the loss of the cold world.

How It Actually Works

Ice arrives before the Ice Age

Thirty-four million years ago, Antarctica crossed into a new condition. Atmospheric carbon dioxide had declined, ocean gateways and circulation were changing, and summer snow on the continent began surviving at a scale that allowed a lasting ice sheet to grow. The date marks the opening of the current long icehouse, although Antarctic ice later expanded and contracted.

The Northern Hemisphere took much longer to acquire comparable continental ice. Mountain glaciers existed earlier, but major glaciation intensified around 2.7 million years ago. By then the positions of continents, ocean circulation and greenhouse-gas levels allowed northern summer snow to accumulate across broad land areas. Ice-rafted debris became more abundant in North Atlantic sediments as glaciers carried rock to sea and icebergs released it offshore.

The formal Pleistocene begins at 2.58 million years ago. Nothing magical happened on the boundary itself. Geological periods are lines drawn through continuous change, placed where a recognisable global signal lets different records be compared. The new epoch captures the interval in which repeated Northern Hemisphere glaciations became the dominant environmental rhythm for many large land animals, including early humans.

It also sits inside the Quaternary Period, which continues through the Holocene. Quaternary therefore names the whole recent period; Pleistocene names its long glacially restless first epoch.

At first the ice sheets were generally smaller and more frequent than the later giants. The strongest beat followed Earth's changing axial tilt, near 41,000 years. Marine sediments preserve the alternation as shifts in oxygen isotopes within the shells of tiny organisms. When more water was locked in land ice, the ocean became enriched in the heavier oxygen isotope. Layer after layer turned changing ice volume into a long, indirect clock.

A 41,000-year world

Early Pleistocene landscapes changed repeatedly without becoming empty between cycles. Grasslands expanded and contracted. Forests shifted latitude and altitude. Rivers cut terraces as sea level and sediment supply changed. Animals moved between Africa and Eurasia when routes and habitats allowed. Some lineages adapted, some split, and many disappeared long before the best-known late Ice Age fauna existed.

Human populations were already part of that movement. Members of the genus *Homo* left Africa and reached Eurasia early in the Pleistocene. Their tools show continuity and local change across vast distances, but a stone artefact does not identify one neat migrating people. Populations could expand, pause, vanish and be replaced while similar technologies persisted. The Ice Age repeatedly edited the human range before *Homo sapiens* appeared.

The 41,000-year rhythm did not mean each glacial cycle copied the last. Ice growth depended on the geography left behind, the state of the oceans and atmosphere, and whether old regolith or sediment beneath an ice sheet made the bed easier to slide over. A cycle could begin under similar orbital geometry and still produce a different ice volume. The planet carried memory forward.

Between about 1.25 million and 750,000 years ago, that memory changed the rhythm. Glacials grew longer and larger, with a strong average spacing

near 100,000 years. The shift is called the Mid-Pleistocene Transition. It did not coincide with a new orbital cycle powerful enough to account for it. Explanations include erosion of soft material beneath ice sheets, allowing them to become thicker on harder bedrock; changes in carbon dioxide; altered ocean circulation; and nonlinear ice-sheet thresholds. Each may contribute. None has closed the case.

The later Pleistocene therefore developed a saw-toothed pattern. Ice sheets often grew over many tens of thousands of years, then collapsed more rapidly. The long build and faster release reflected the interaction of orbital pacing with greenhouse gases, reflectivity and ocean heat transport. The resulting world spent more time in cold conditions than in warm peaks like the present interglacial. Larger ice sheets also changed their own surroundings. Their height cooled the air above them, their margins produced meltwater and dust, and their weight altered the bed. An ice sheet was both a response to climate and a new geographical feature capable of modifying climate.

The last warm interval

Around 129,000 years ago, the previous interglacial reached full warmth. Forests occupied regions later covered by tundra or ice. Sea level stood higher than today because oceans were warmer and global land ice was smaller, although the exact height and contributions of Greenland and Antarctica remain active research questions. Hippopotamuses ranged in parts of Britain during the wider warm interval, a useful correction to the idea that northern Europe has one natural climate.

The warmth did not fail because an interglacial has a fixed expiry date. Orbital geometry gradually reduced summer sunlight at high northern latitudes. Snow survived in favourable places, ice expanded, reflectivity rose and greenhouse-gas concentrations declined. By about 115,000 years ago, the last glacial period was under way. Sea level began a long, uneven fall. Britain changed from warm woodland towards colder open environments, but the transition took many millennia and included reversals. Elsewhere, rainfall mattered more than frost. North African lakes, Arabian corridors and

Asian monsoons expanded or contracted as orbital seasonality and glacial boundary conditions changed, reshaping routes far from any ice sheet.

Its descent was irregular. Marine Isotope Stages divide the record into numbered warm and cold intervals, with even numbers generally glacial and odd numbers generally warmer. Those global labels conceal violent regional variation. Greenland ice cores from the middle of the last glacial period record repeated abrupt warmings followed by slower cooling, known as Dansgaard-Oeschger events. North Atlantic sediments contain layers of debris delivered by armadas of icebergs during some cold episodes, called Heinrich events. Ocean circulation, ice-sheet instability and atmospheric change were coupled tightly enough for one region to shift while the slow orbital background barely moved.

Living through this world required flexibility across generations. A productive valley could dry. Open steppe could become shrubland. A coast could move outward as sea level fell, then become a route for people and animals. Glaciers could block one pass and create another margin to follow. Archaeological sites capture brief occupations within a landscape whose usable places changed repeatedly.

Several human worlds meet

Neanderthals had occupied Europe and western Asia across earlier cold and warm cycles. They used fire, hafted tools, hunted large animals, gathered plants and maintained social groups capable of caring for injured members. Their range shifted with climate, but they were not restricted to snow or one dietary strategy. The label cold-adapted is useful for parts of their anatomy and history, then dangerous if it turns them into specialists unable to live outside tundra.

Farther east, Denisovans left a thinner skeletal record but a strong genetic one. A finger bone and teeth from Denisova Cave helped reveal a population related to Neanderthals that contributed ancestry to later humans. Other finds and genomes have widened their known geography. The mismatch between a few fossils and a large genetic inheritance is a

reminder that the human fossil record is not a census.

Homo sapiens expanded from Africa in several movements. By the later part of the last glacial period, modern human populations occupied large parts of Eurasia and Sahul. Some early expansions ended without supplying much ancestry to later people. Others met Neanderthals and Denisovans and interbred with them. Genetic exchange means the clean branching diagram taught in older accounts must be redrawn as branches that sometimes rejoin.

In Europe, improved radiocarbon chronologies place the disappearance of the last securely dated Neanderthal populations around 40,000 years ago, with overlap between Neanderthals and incoming modern humans lasting millennia and varying by region. Climate was unstable during this interval, but no cold event singles itself out as a universal executioner. Neanderthal populations were small and fragmented. Modern human networks may sometimes have been larger or more connected. Competition, disease, assimilation and demographic chance remain possible contributors.

The archaeological sequence does not support a single march of progress. Tool traditions overlap. Innovations appear, spread unevenly and sometimes disappear. Needles, tailored clothing, projectile systems, ornaments, pigments and art become visible in different places and periods, while organic technologies are badly underpreserved. A bone awl may survive where an entire hide shelter does not. What looks like a sudden cognitive leap can partly reflect what the ground kept.

By the time ice sheets reached their last maximum, human populations had survived many reversals. Their success was conditional. Northern Europe was largely abandoned where ice or polar desert removed food. Populations concentrated in refugial regions, river corridors, coasts and productive steppe. Elsewhere, from Africa to southern Asia, the glacial world was experienced through drought, changing monsoons and shifting shorelines rather than nearby continental ice. A lowered coast could expose new foraging ground while moving fresh water farther from the old shore. A colder dry phase could open grassland to grazing prey while reducing plants

used in another season. Human flexibility worked by solving local combinations, not by carrying one Ice Age kit everywhere.

The last maximum

From about 26,500 years ago, nearly all major ice sheets occupied positions associated with the global Last Glacial Maximum. In North America, Laurentide ice merged or met with western ice over parts of the continent. Northern Europe lay beneath the Fennoscandian and British-Irish ice sheets. Greenland and Antarctica held more ice than today, while glaciers enlarged across high mountains in both hemispheres.

The term maximum needs care. Different sectors reached their farthest positions at different times, and some mountain glaciers had local maxima outside the global interval. The label describes a broad peak in global ice volume, not one date on which every glacier stopped in formation for a photograph.

Global mean cooling near 6 degrees Celsius transformed regional climates. Over and beside the northern ice sheets, cooling was much larger. Permafrost spread. Wind carried dust across sparsely vegetated ground. The Sahara and other dry regions changed extent as rainfall belts shifted. Tropical forests contracted or altered composition in many areas, while mountain vegetation descended. In some basins, lakes grew because evaporation fell or storm tracks moved; elsewhere lakes shrank. There was no universal glacial weather.

About 125 to 130 metres of sea-level fall exposed enormous shelves. The English Channel was land crossed by river systems. The Beringian region joined Asia and North America, although glaciers and ecology governed passage beyond it. The Sunda Shelf created a larger Southeast Asian landmass, while people reaching Sahul still had to cross open water. Reduced ocean area and altered coastlines changed tides, estuaries and marine productivity as well as terrestrial routes.

Across northern Eurasia and Beringia, cold open ecosystems supported large herbivores in regionally different mixtures. Mammoths used tusks and

trunks to uncover vegetation beneath snow. Reindeer migrated. Horses and bison grazed. Cave lions and wolves hunted. Humans did too, but archaeological survival gives some places far more detail than others. A decorated cave is an archive, not an average dwelling. Most lives happened in camps whose wood, skin, fibre and food waste vanished. Where occupation remained possible, groups had to anticipate seasonal scarcity. Reliable fuel, food that could be carried or preserved, repeated knowledge of migration routes and access to good stone could determine whether a site was a useful base or a brief stop. Cold was experienced through logistics.

Ice sheets were active bodies throughout the maximum. Snow accumulated in cold interiors. Ice flowed towards margins, deforming under its own weight and sliding where water or sediment allowed. Outlet glaciers calved. Margins advanced, paused or retreated. The apparent stillness of an ice map compresses a moving mass and thousands of local balances.

The release

Around 19,000 to 20,000 years ago, rising northern summer sunlight helped start the long release from maximum ice. Melting exposed darker ground, atmospheric carbon dioxide rose, and ocean circulation redistributed heat. The ice sheets did not retreat at a fixed speed. Margins stalled, readvanced and broke apart. Meltwater collected against ice dams, escaped through new outlets and sometimes entered the ocean in abrupt pulses.

Sea level responded with delay because so much ice had to move from land to ocean. Over thousands of years it rose more than a hundred metres. Each rise redrew the coastal world. Flat shelf landscapes were narrowed, divided and drowned. People could retreat upslope or inland, but camps, hunting grounds and routes disappeared beneath water. Much of the archaeology of coastal Ice Age life is therefore inaccessible on the seabed. Inland, retreating ice left unstable drainage. Proglacial lakes occupied low ground against the ice margin, and their outlets could switch as a margin thinned. Rivers carried enormous sediment loads, building floodplains and deltas that later communities inherited. The thaw created land at the same

time as it removed it.

Around 14,700 years ago, Greenland and much of the North Atlantic region warmed sharply at the start of the Bølling-Allerød interval. Ice retreated and vegetation changed quickly. Then, around 12,900 years ago, the Younger Dryas reversed much of the regional warming. Greenland temperatures fell, glaciers advanced in some mountains and tundra returned across parts of Europe.

The event's name comes from *Dryas octopetala*, a small Arctic-alpine flower whose pollen and leaves reappeared in European sediments as cold conditions returned. Its cause has often been reduced to meltwater shutting down Atlantic circulation. Freshwater input and weakened overturning remain leading parts of many explanations, but modelling and records indicate a combination of ocean, atmospheric and radiative changes. The impact was strongest around the North Atlantic. The Southern Hemisphere did not enter a matching deep freeze.

At 11,700 years ago, rapid warming in Greenland marks the formal start of the Holocene. That boundary is crisp in the timescale and messy on the ground. Remnant ice sheets persisted. Seas continued rising for millennia. Forests expanded along species-specific routes. Cold-adapted animals withdrew northward or uphill. Human communities changed mobility and diet in different ways, then in some regions began the long transition towards cultivation and herding. Britain remained connected to continental Europe for part of this early interval before rising water finally divided the lowlands. The drowning was progressive, not a single morning on which an island appeared. Across the drowning, successive generations inherited coastlines and routes different from earlier ones.

What made it through

The end-Pleistocene extinction pattern was already under way before the Holocene line and continued after it in isolated populations. Woolly rhinoceroses disappeared before the final boundary. Mammoths vanished from most mainland regions but survived on islands. On Wrangel Island, a population lasted until roughly 4,000 years ago. Their late survival shows that suitable local conditions could postpone extinction, while isolation and small population size created new vulnerabilities.

The continental sequence differed sharply. Many Australian giants had gone tens of thousands of years earlier. Eurasian losses were spread across a long period of climatic and human change. The American pulse was later and more concentrated near the Pleistocene's close. These timings matter because a cause that fits one continent may fail on another. Human arrival, population growth, changing vegetation and abrupt climate must be compared against local extinction dates rather than averaged into one global moment.

Other Ice Age animals reached the present. Reindeer, musk oxen, wolves, brown bears, saiga, bison and horses survived in altered distributions, though particular species or regional lineages were lost. In the Americas, horses and mammoths vanished while bison persisted. In Eurasia, reindeer survived but woolly rhinoceroses did not. Such contrasts resist a single body-size rule. Plants recolonised from multiple refugia. Human genomes retained Neanderthal and Denisovan ancestry. Glacial erosion, loess, rebound and drowned shelves preserved the physical inheritance.

The survivors did not restore the old world. A community is not stored inside one species, ready to unfold when climate returns. The mammoth steppe dissolved because its climate, plants, herbivores and geography no longer coincided. The Holocene inherited components, fragments and absences. It became a new assembly built from what remained.

How we know

The Ice Age is reconstructed from records that fail in different ways. Moraines, striations, erratic boulders, till and raised shorelines map former ice and water, but later glaciers can erase earlier limits. Marine sediments preserve oxygen-isotope cycles and ice-rafted debris across millions of years, although isotopes combine temperature and ice-volume effects. Ice cores trap old air, dust and isotopic temperature signals in annual or compressed layers, with dating precision declining deeper in the core.

Pollen, beetles, bones, plant remains, cave deposits and lake sediments reconstruct local environments. Radiocarbon dates recent organic material but requires calibration and ends near the practical limits of the method. Ancient DNA identifies ancestry and former presence, while surviving best in cold settings and remaining vulnerable to contamination and transport within sediment. Archaeology records human behaviour through a severe filter that favours stone, bone and caves over fibre, wood and ordinary camps.

Confidence comes from convergence. No single core, fossil or model supplies the story. Uncertainty remains largest where records are sparse, dates overlap, causes interact or a local sequence is made to represent a continent. The broad cycles are secure. The exact route by which each population met them often remains open.

What People Get Wrong

“The whole planet was buried under ice”

The image survives because an ice sheet is visually decisive. A map coloured white over Canada and northern Europe becomes, in memory, a white Earth. Films add blizzards where the map leaves space.

At the Last Glacial Maximum, continental ice covered large parts of northern North America and Europe and expanded in many mountain ranges. Most land and ocean remained uncovered. Sea ice expanded seasonally, but floating sea ice should not be confused with the land ice that lowered global

sea level. The global mean was about 6 degrees Celsius colder than the pre-industrial mean, but the change was uneven. Some high-latitude regions were far colder; many tropical regions cooled less. Outside the ice margins, dryness, wind, altered monsoons and lower carbon dioxide could matter as much as frost.

This was not a minor freeze. Sea level fell by roughly 125 to 130 metres, permafrost spread and ecosystems shifted across continents. White coverage is therefore a poor measure of climate impact. A region can be transformed without a glacier reaching it. The Ice Age was planetary because circulation, water storage and habitats changed, not because the whole globe became an ice rink.

“There was one Ice Age”

The singular is convenient enough to become a trap. Museum galleries and school timelines usually begin with familiar mammoths and end when the ice retreats, compressing 2.58 million years into one cold episode.

The Pleistocene contained many glacials and interglacials. Ice sheets grew, shrank and returned. Their rhythm changed through time, from cycles strongly associated with axial tilt to longer, larger later cycles. The Last Glacial Maximum was the peak of the most recent glacial cycle, not the peak of every glacier at one instant and not the whole Pleistocene.

Earth has also experienced much older episodes of widespread ice. The Quaternary story is merely the one that shaped modern landscapes, living species and human populations most directly. Even now, permanent ice remains on Greenland and Antarctica, so geologists can describe Earth as still occupying a long icehouse state.

Repeated change selects for different things from one long emergency. A refuge that works during one glacial may fail in the next interglacial. The defining pressure was not endless cold. It was the repeated movement of the conditions organisms needed.

“The coldest winters built the ice sheets”

Winter is when snow falls, so the intuition appears sound. Yet an enormous winter snowfall followed by a warm summer leaves no long-term ice. Ice sheets require a positive balance across years: accumulation must exceed melting, sublimation, flow and iceberg loss.

Cool summers at high northern latitudes therefore had exceptional leverage. Changes in orbital geometry redistributed seasonal sunlight. When summer energy weakened, snow could survive, reflect more sunlight and accumulate. Strong snowfall still mattered because a dry region cannot build ice from cold alone. The favourable combination was enough moisture to add snow and summers cool enough not to remove it.

This explains why glaciation cannot be read from one weather record. A harsher winter may occur within a climate whose summers still destroy the snowpack. An ice sheet can also keep growing in its cold interior while losing mass rapidly at a warm or marine margin.

The same error reaches beyond ice. Annual averages can hide the season that controls a system. A small change at the limiting moment can matter more than a dramatic change at the wrong one.

“Mammoths lived in a frozen desert”

Mammoths are drawn among snowdrifts because snow communicates cold at once. The picture leaves an obvious problem: several tonnes of herbivore need a great deal of food.

The mammoth steppe was a broad family of cold, dry, open ecosystems rich in grasses, sedges, flowering herbs and low shrubs. Shallow snow could leave forage accessible through winter. Long summer daylight and mineral-rich dust supported productivity in some areas. Large herbivores may have helped keep ground open through grazing and trampling, but moisture, soils and climate constrained the result.

It was not uniform. Ancient environmental DNA from Arctic sediments shows regional mixtures of plants and changing communities across the last

50,000 years. The evidence also comes from selected cold regions and cannot represent every mammoth habitat. Mammoths lived from western Europe through northern Asia and North America across different climates and times.

Cold does not mean biologically empty. The lost world supported high animal biomass through a combination that has largely disappeared. Warming removed that combination rather than revealing a timeless green normal underneath it.

“Humans survived because they were the smartest species”

The claim flatters the survivor and explains almost nothing. Intelligence has no single archaeological measure, and every extinct human population is then defined as less intelligent by the fact of its extinction. Brains do not fossilise as behaviour, while the tools and shelters that would reveal planning survive selectively. Absence can reflect decay as easily as incapacity.

Neanderthals made sophisticated tools, used fire, hunted difficult prey and cared for injured people. Denisovans left sparse fossils but a wide genetic legacy. Modern human populations also vanished repeatedly. The lineages that endured did so through particular combinations of demography, mobility, technology, diet and social connection. Wider exchange networks may have buffered some groups, but the evidence varies and does not establish one universal cognitive advantage.

Interbreeding further damages the contest. Neanderthal and Denisovan ancestry survives in living people. Some groups disappeared as distinct populations while contributing to others. Survival is therefore not a podium with one clean winner.

Replacing destiny with conditions changes the human story. Culture made humans unusually flexible, but flexibility had to be maintained by people, knowledge and contact. A clever individual inside an isolated collapsing population is still inside an isolated collapsing population.

“People killed every giant animal”

Some accounts turn late Pleistocene extinction into a single hunting spree. Others react by removing humans and blaming climate. Both offer the pleasure of one culprit.

Humans clearly hunted some large animals, and broad geographical patterns give human expansion a major causal role. Large, slow-breeding species can decline when even a small extra share of adults is killed. Yet kill sites are uneven, extinction dates differ, and climate changed habitat, water and population connectivity. Australia, Eurasia and the Americas did not lose their megafauna at one time or under identical conditions.

The strongest account is interaction with species-specific weighting. Human pressure may be decisive where prey had little prior exposure to people or populations were already small. Habitat change may dominate another phase. Disease, fire and competition are harder to test and should not be inserted merely to complete a list. The balance must be established locally. That requires population-level dates, habitat evidence and an estimate of how much extra mortality a slow breeder could absorb.

The false trial is “humans or climate”. A population reduced by disappearing habitat can be finished by hunting that would once have been sustainable. Causes combine inside a demographic process.

“The Ice Age ended, so the ice story is over”

The Pleistocene ended 11,700 years ago, and the last glacial period had already given way to rapid warming. In that chronological sense, the familiar Ice Age is over. The Holocene is the current warm interval within the Quaternary, not a return to the world that existed before northern glaciation began.

Its machinery and inheritance remain. Greenland and Antarctica still hold permanent ice. Formerly glaciated land continues to rebound. Rivers, lakes, soils and coastlines occupy forms made by ice and meltwater. Living populations carry genetic divisions, bottlenecks and admixture formed

during glacial retreat and recolonisation. Britain is an island because postglacial seas drowned the lowlands that connected it to Europe.

The present climate is not another ordinary stage of orbital deglaciation, and ancient cycles do not explain modern industrial warming. Orbital forcing evolves over thousands of years and current greenhouse forcing has a different source and pace. Treating every climate change as interchangeable destroys causal understanding.

An ending can leave an operating inheritance. The ice sheets retreated, but they did not restore the previous map. We live on the rearranged ground among the selected survivors.

Use It

Separate the pacemaker from the amplifier

Orbital geometry paced glacial cycles, but the orbit did not manufacture kilometres of ice directly. Snow and ice reflected sunlight. Greenhouse gases changed. Oceans moved heat and carbon. Vegetation and dust responded. The first disturbance and the largest source of change were not identical.

Carry that distinction into any explanation involving feedback. Ask what altered the timing, what enlarged the response and what stored memory between episodes. A regular trigger can produce irregular outcomes when the system begins each cycle in a different condition. A weak initial shove can matter because it crosses a threshold, while a dramatic variable may be a response that later becomes a cause in its own right.

This lens also stops false either-or arguments. Carbon dioxide can amplify an orbitally paced transition without being the first variable to move. Ice can be an outcome of cooling and a driver of further cooling. Causal roles change through the sequence. Naming one original trigger does not dismiss the machinery that turns it into a planetary event.

Ask what moved across the map

A global temperature number is necessary and incomplete. During the Last Glacial Maximum, the mean cooling was about 6 degrees Celsius, yet organisms experienced moving snowlines, rainfall belts, coasts, rivers, forests and prey. The practical event was a relocation of usable habitat.

When assessing past environmental change, redraw the map. Which shelves emerged as sea level fell? Which routes were blocked by ice? Which valley remained wet? Which mountain created a cool refuge during warming? The same average change can connect populations in one region and divide them in another.

This approach prevents geography from becoming passive scenery. Doggerland existed because ocean water was stored on land. Beringia was useful only while its ecology and onward routes were workable. A coastline that shifts fifty kilometres can alter fresh water, food and travel even when the people involved never see an ice sheet. Temperature is an input. Access is the lived consequence.

Find the controlling season

Annual averages hide the condition that decides whether a process continues. Ice sheets need snow, but cool summers determine whether enough survives. A cold winter followed by strong melt can produce less lasting ice than a milder winter followed by a cool summer. In mammoth-steppe habitats, winter snow depth could matter more to grazers than the lowest temperature because deep snow sealed food away.

The same discipline improves palaeoclimate reading. Ask whether an orbital change altered summer sunlight, monsoon seasonality or winter storm tracks rather than treating one annual mean as the mechanism. Plants respond to growing seasons and frost timing. Rivers respond to the order of snow storage and melt. Animals meet the week of scarcity, not a yearly average printed in a table.

A limiting season is not always the same across regions or species. The

value of the lens lies in finding it rather than assuming it. The decisive variable is often the one that acts at the bottleneck in time.

Find the refuge and the route

A species range on a map can make survival look like a property of the organism. The Ice Age shows that persistence often belongs to a relationship among organism, refuge and corridor. A population must retain somewhere suitable, then reach new habitat when conditions move.

Look for the refuge at the right scale. Southern Europe may be a refuge in a continental summary, while one sheltered slope, spring-fed valley or coastal plain may be the usable area for a particular population. Cold-adapted species need refuges during warmth as temperate species do during glacials. Safe ground can therefore switch direction between cycles.

Then examine connection. Is the new habitat continuous, or separated by forest, water, ice or unsuitable soil? Does the species disperse quickly enough? Can its prey, host or pollinator move with it? A corridor may also carry competitors, predators and disease. Connectivity is not automatically beneficial, but without some route a shrinking refuge becomes an island. Survival planning begins with the map of possible continuity, not a list of admirable adaptations.

Count survival at more than one level

The word survived can conceal large losses. A species may remain while most of its populations disappear. A population may end as a distinct group while some ancestry continues through interbreeding. A vegetation type may return after ice retreat but with a different combination of species. A broad modern range may descend from a narrow refugium and carry reduced genetic diversity.

Use at least four levels: species, population, community and landscape. Woolly mammoths survived the Pleistocene on isolated islands, but the mammoth-steppe community did not. Neanderthals disappeared as a distinct human population, while Neanderthal ancestry remains in living

people. Reindeer survived, though their range and ecological neighbours changed.

This prevents survival from becoming a yes-or-no scoreboard. Persistence has quality, distribution and inheritance. A line on a species list can stay unchanged while resilience, abundance and connectivity deteriorate. Conversely, a named population can disappear without every biological contribution being erased. The Ice Age record becomes clearer when continuation and replacement are allowed to coexist.

Reconstruct extinction as a sequence

One-cause verdicts are attractive because they make an ending feel solved. Late Quaternary extinctions instead require a timeline for each population. When did suitable habitat shrink? When did humans arrive or intensify their presence? Was the population already isolated? How quickly did it reproduce? Which additional mortality could it absorb?

Order changes interpretation. Hunting that appears light against a large connected population may be severe after climate fragments the range. Warming can increase plant growth in one season while deeper snow blocks winter forage. An island refuge can protect a remnant from mainland pressures, then expose it to inbreeding, drought or one later disturbance. Extinction often arrives when several tolerable pressures become intolerable in combination.

Demand dates at the same geographical scale as the claim. A continental climate curve cannot by itself explain the last population in one valley. One kill site cannot prove a continent-wide extermination. The useful question is not which side wins, humans or climate. It is which sequence drove numbers below recovery here.

The limits

The Pleistocene is not a controlled experiment repeated with one variable changed. Orbital cycles differed, ice sheets inherited different beds, continents held different species, and human populations expanded through the later cycles. Similar-looking outcomes can therefore arise by different routes. A past correlation between warming and extinction does not assign a modern cause, and a surviving species does not reveal every pressure that its lost populations faced.

The evidence has strong geographical biases. Cold caves, permafrost, lake beds and marine sediments preserve different materials. Europe and North America have been sampled more heavily than many regions of Africa, Asia and the tropics. Environmental DNA is powerful where it survives and where reference genomes permit identification. Fossils favour hard parts and depositional settings. The blank spaces on a map are mixtures of absence, non-preservation and non-discovery.

The cycles also unfolded over thousands of years. They show that climate systems possess feedbacks and thresholds, but they are not direct templates for present decisions. Modern greenhouse forcing has a different cause and a pace that cannot be collapsed into the phrase climate has always changed. Analogy is useful only after the forcing, rate, baseline and geography have been compared.

Finally, survival is not moral achievement. Mammoths were not poorly designed because they vanished, and humans did not earn existence through superior worth. Evolution and extinction have no tribunal. They record which populations reproduced under conditions that happened to arrive.

The one thing to keep

Keep the moving map.

The Ice Age is easiest to remember as temperature: colder air, bigger glaciers, warmer air, retreat. Temperature began the explanation and

geography delivered the consequences. Water left the ocean and became land ice. Shelves emerged. Rivers turned. Rainfall belts shifted. Forests broke apart. Refugia appeared in one cycle and failed in the next. Routes joined populations, then vanished beneath rising seas.

Every animal, plant and human group met that movement at its own speed. Thick fur mattered only where food remained. Intelligence mattered only where people, knowledge and cooperation persisted. A species could endure in one hidden pocket while disappearing across a continent. A population could vanish and leave genes behind. The final thaw could be as destructive to a cold specialist as the preceding freeze was to a warm one.

That should permanently alter the picture. The survivors of large environmental change are not always the strongest occupants of the old world. They are the populations that remain connected to a workable next place, through movement, refuge, flexibility or inheritance. Look at any modern distribution and ask where it waited, how it returned and what never came back with it.

The world beneath you is not the stage on which the Ice Age happened. It is one of the outcomes.

Terms

Ice age. A long interval when permanent ice sheets exist on Earth, often containing many colder glacials and warmer interglacials. In popular use, it usually means the Pleistocene.

Icehouse. A planetary climate state with substantial permanent polar ice. It contrasts with greenhouse states that lack large enduring ice sheets, though temperatures still vary within either state.

Quaternary. The geological period from 2.58 million years ago to the present. It contains the Pleistocene and Holocene and is marked by repeated Northern Hemisphere glaciation. Its name is historical, not a claim about four repeating units.

Pleistocene. The epoch from 2.58 million to 11,700 years ago. It contains most of the familiar glacial cycles, Ice Age megafauna and much of human evolution. The name arose from nineteenth-century fossil classification.

Holocene. The current geological epoch, beginning 11,700 years ago after abrupt deglacial warming. It is a warm interval within the continuing Quaternary and long Cenozoic icehouse. Its formal lower boundary is defined from Greenland ice.

Glacial. A colder interval during an ice age when continental ice sheets and mountain glaciers generally expand and global sea level falls. Regional timing and severity need not match.

Interglacial. A warmer interval between glacials when ice sheets retreat, sea level rises and ecosystems reorganise. The Holocene and the last interglacial are separate examples, with different orbital geometry, ice extent and regional climates.

Stadial and interstadial. Relatively cold and warm phases within a broader glacial or interglacial sequence. Greenland records show sharp interstadials during the last glacial period without a full return to interglacial conditions; regional timing may differ.

Younger Dryas. A severe return to cold around the North Atlantic from roughly 12,900 to 11,700 years ago, interrupting deglacial warming. It was strongest in the north and was not a second worldwide glacial maximum.

Last Glacial Maximum. The broad global interval, roughly 26,500 to 19,000 or 20,000 years ago, when nearly all major ice sheets occupied positions associated with maximum ice volume.

Marine Isotope Stage. A numbered interval defined mainly from oxygen-isotope changes in marine sediments. Even numbers are generally colder and odd numbers warmer, with important local exceptions.

Milankovitch cycles. The combined orbital variations in eccentricity, obliquity and precession that redistribute sunlight by latitude and season and pace glacial-interglacial change over long timescales. They are named

after Milutin Milankovitch, who calculated their climatic effects.

Eccentricity. Variation in the shape of Earth's orbit from more circular to slightly more elliptical. Its direct forcing is weak, but it alters the strength of precession.

Obliquity. The changing tilt of Earth's axis, with a rhythm near 41,000 years. Greater tilt strengthens seasonal contrast, especially at high latitudes, and affects summer melting.

Precession. The wobble of Earth's axis and rotation of its orbital orientation. It changes which hemisphere's summer occurs nearer the Sun and modifies seasonal sunlight on rhythms near 19,000 to 23,000 years.

Insolation. Incoming solar radiation received at a place and season. Glacial theory pays particular attention to summer insolation at high northern latitudes because it controls snow survival.

Albedo. The fraction of incoming sunlight a surface reflects. Bright snow and ice have high albedo; darker land and open water absorb more energy and reinforce warming.

Climate feedback. A response that changes the original disturbance. Ice-albedo and greenhouse-gas feedbacks amplified orbital pacing, while the size and timing of each response varied across cycles.

Ice sheet. A land-based mass of glacial ice covering more than 50,000 square kilometres. Ice sheets flow under their own weight and lower sea level as they grow.

Glacier. A persistent body of land ice formed from accumulated snow and moving under gravity. Mountain glaciers are smaller than ice sheets but preserve local climate and landform evidence.

Firn. Old compacted snow whose grains have partly fused but have not yet become solid glacial ice. It records the transition from annual accumulation to long-term storage.

Mass balance. The difference between ice gained through snowfall and

refreezing and ice lost through melting, sublimation and calving. Positive balance grows ice; negative balance shrinks it.

Moraine. A ridge or deposit of rock and sediment carried and left by a glacier. Moraines can mark former margins, pauses or readvances, though later erosion can obscure them.

Till. Unsorted sediment deposited directly by glacier ice, ranging from clay to boulders. Its mixed texture distinguishes it from material sorted by rivers, waves or wind.

Loess. Fine windblown sediment, often sourced from glacial grinding and exposed floodplains. Thick loess deposits preserve climate records and have produced fertile soils after weathering.

Permafrost. Ground remaining at or below freezing for at least two consecutive years. It can preserve organic remains, restrict drainage and roots, and reshape slopes when it thaws.

Beringia. The exposed and adjacent land region that connected northeast Asia and northwest North America during low sea level. It was habitat and homeland, not merely a bridge.

Refugium. An area where a population persists while conditions become unsuitable across much of its range. Refugia are species-specific, temporary and often genetically important sources of recolonisation.

Megafauna. Large animals, defined differently among studies and regions. In Ice Age discussion it usually refers to large mammals whose late Quaternary losses were unusually severe.

Mammoth steppe. A varied belt of cold, dry, open vegetation across northern Eurasia and Beringia that supported mammoths and other grazers and has no exact modern equivalent.

Go Deeper

Jamie Woodward, *The Ice Age: A Very Short Introduction*

Oxford University Press, 2014. Start here for the compact physical overview. Woodward moves from the discovery of former glaciation through orbital cycles, ice sheets, landforms, sea level, palaeoclimate evidence and human occupation without treating the subject as a mammoth catalogue. The format is close to this book's promise but more geological in emphasis, so it is useful for checking how moraines, sediments and isotope records become a history of vanished ice. It also keeps older glaciations in view, preventing the Pleistocene from being mistaken for Earth's only frozen interval. The compression is brisk and some diagrams deserve slower attention than the page count suggests.

Richard B. Alley, *The Two-Mile Time Machine*

Princeton University Press, 2000. Read this for the evidence inside ice and for abrupt climate change as a scientific problem. Alley helped study the Greenland ice cores and explains how snowfall becomes a layered archive containing isotopes, dust, chemistry and trapped air. The book follows the practical difficulty of drilling, dating and interpreting cores, then shows why sharp shifts in Greenland changed the old picture of gradual deglaciation. It is written for general readers and remains unusually clear. Its discussion predates two decades of later cores, models and dating refinements, so use it for method and intellectual discovery rather than as the final word on every mechanism.

Anthony J. Stuart, *Vanished Giants: The Lost World of the Ice Age*

University of Chicago Press, 2021. Read this for the animals, their changing ranges and the extinction argument. Stuart is a palaeontologist who treats mammoths, woolly rhinoceroses, giant deer, cave lions and other large mammals as populations moving through particular environments rather than as a poster of doomed monsters. The book's geographical breadth makes the uneven timing of losses hard to ignore and gives due weight to climate, humans and interaction. It is more detailed than a one-hour survey and strongest on Eurasia, where Stuart's own work is concentrated. That setting makes it a major interpretation, not a universal template to be applied without local dates.

Rebecca Wragg Sykes, *Kindred: Neanderthal Life, Love, Death and Art*

Bloomsbury Sigma, 2020. Read this for the inhabited human world. Wragg Sykes reconstructs Neanderthal bodies, food, tools, movement, social care and symbolic behaviour from archaeological evidence while showing how much depends on preservation and inference. It is the best antidote to the lumbering cold-specialist caricature and to the opposite temptation to remake Neanderthals as modern people in different clothes. The book is long, richly detailed and openly interpretive. Use it alongside current dating and genomic research, since the evidence changes quickly, but begin here when stone tools and bones need to become lives without invented scenes.

Notes and Sources

Scope, dates and terminology

The book uses Ice Age in the familiar sense of the Quaternary glacial world, centred on the Pleistocene, while distinguishing that interval from the longer Cenozoic icehouse. The formal Pleistocene base is 2.58 million years ago and the Holocene base is 11,700 years ago in the current International Chronostratigraphic Chart. Geological boundaries organise records; they do not imply instantaneous global environmental change.

The statement that large Antarctic ice formed around 34 million years ago and major Northern Hemisphere glaciation intensified around 2.7 million years ago follows the Cenozoic synthesis of Zachos and colleagues and later Quaternary reviews. The exact onset and stability of ice differed among sectors and through time. Ice age and icehouse are used as broad state descriptions, not as claims that ice sheets were fixed.

Orbital pacing and glacial rhythm

Hays, Imbrie and Shackleton established the classic empirical case that orbital variations paced late Pleistocene climate. Eccentricity, obliquity and precession redistribute sunlight by latitude and season rather than creating a large change in annual global solar energy. The emphasis on high northern summer sunlight follows orbital ice-sheet theory and the geography of northern land masses. Snowfall remains necessary; cool summers alone cannot grow ice in a region with too little accumulation.

The marine oxygen-isotope chronology follows the LR04 stack of Lisiecki and Raymo. Benthic oxygen isotopes respond to both global ice volume and deep-ocean temperature, so they are not a direct ice gauge without interpretation. The early 41,000-year rhythm, later near-100,000-year cycles and timing of the Mid-Pleistocene Transition follow Herbert's 2023 review. Proposed mechanisms include regolith removal, ice-sheet thresholds, ocean circulation and carbon-cycle change. The manuscript retains the unresolved status rather than selecting one mechanism.

Feedbacks and greenhouse gases

The EPICA Dome C core and later carbon-dioxide measurements extend Antarctic climate and greenhouse-gas evidence across multiple glacial cycles. They show close coupling among Antarctic temperature, atmospheric carbon dioxide and glacial state. The manuscript does not treat carbon dioxide as a passive thermometer or as the sole initial trigger. Orbital changes, ice, oceans and carbon interact, and the sequence differs by timescale and region.

The distinction between pacemaker and amplifier is explanatory rather than a quotation from one paper. Hays and colleagues support orbital pacing; EPICA and Lüthi and colleagues support greenhouse-gas covariation; Herbert reviews the nonlinear system and the continuing 100,000-year problem.

Last Glacial Maximum scale

Clark and colleagues define the broad global Last Glacial Maximum interval used here: ice sheets grew towards maximum positions from about 33,000 to 26,500 years ago, and nearly all were near LGM positions from 26,500 to 19,000 or 20,000 years ago. Local glacier maxima differ, which is why the text avoids one universal peak date.

Tierney and colleagues estimated global mean LGM cooling at 6.1 degrees Celsius relative to the pre-industrial climate, with a 95 per cent interval of 5.7 to 6.5 degrees. The body rounds this to about 6 degrees and stresses spatial variation. Lambeck and colleagues reconstructed global sea level and ice volume from roughly one thousand observations, with the low stand around 130 metres below present in their preferred curve. The body uses roughly 125 to 130 metres because estimates vary with method, correction and reference level.

Ice sheets, landforms and drowned landscapes

The treatment of glacier formation, flow, mass balance, moraines, till, loess and isostatic rebound follows Woodward's geological synthesis. The Great Lakes sentence is deliberately bounded: repeated ice modified and deepened basins controlled by older bedrock rather than cutting an entirely new landscape from uniform rock.

Gaffney, Fitch and Smith provide the main source for Doggerland as a lived landscape beneath the North Sea. Seabed finds and geophysical mapping support rivers, wetlands, fauna and human presence. Its drowning was progressive as sea level rose. Beringia is treated in the same way: a broad ecological region that sometimes connected populations, not a permanent road or a single migration event. The book avoids a fixed width or a definitive first-entry chronology for the Americas because both routes and dates continue to be refined.

Mammoth-steppe ecology

Stuart supplies the broad palaeontological account of large mammals, ranges and extinctions. Wang and colleagues analysed 535 permafrost and lake-sediment samples from 74 circumpolar sites spanning the past 50,000 years. Their results support a widespread steppe-tundra flora during the LGM, later regional divergence, a negative association between wetter conditions and grazing-animal diversity in the sampled Arctic regions, and late mammoth persistence in northern Siberia.

That study is the most setting-specific evidence retained in the book. It is powerful for the circumpolar Arctic and cannot establish a global extinction cause. Its abstract also reports humans as a minor factor in animal distributions within that dataset. The manuscript does not generalise that result beyond its spatial and analytical scope because broader archaeological and macroecological evidence gives humans a major role in many extinctions.

The role of large herbivores in maintaining open vegetation remains partly contested. Grazing, trampling and nutrient movement can affect plant

communities, but climate, moisture and soils constrain those effects. The text therefore says herbivores may have helped maintain some open conditions and rejects the stronger claim that mammoths created a uniform biome.

Refugia, movement and genetic inheritance

Stewart, Lister, Barnes and Dalén review the individualistic nature of Quaternary refugia. Refugia can differ among species in location, size and timing; cold-adapted taxa also require refugia during warm periods. Hewitt synthesises the genetic effects of contraction, isolation, expansion and secondary contact. The text does not turn the familiar southern-European tree-refugia model into a universal rule and allows northern, high-altitude, coastal and small local refugia.

Lorenzen and colleagues combined ancient DNA, species-distribution models and the human fossil record for six large mammal taxa. They found climate to be a major population driver while responses to climate, habitat and humans differed by species. This supports the manuscript's refusal to move whole communities as one unit or to predict extinction from body size alone.

Human populations

The human account is intentionally bounded. Wragg Sykes supplies the archaeological synthesis of Neanderthal bodies, technology, subsistence, care and symbolic evidence. Higham and colleagues provide the improved radiocarbon chronology placing Neanderthal disappearance at different times across Europe around 40,000 years ago and allowing a period of overlap with modern humans. This chronology has continued to be tested and refined, so the manuscript avoids a single last-Neanderthal date.

Meyer and colleagues provide the high-coverage Denisovan genome; Prüfer and colleagues provide a high-coverage Neanderthal genome from Vindija Cave. Together with the wider genomic literature, these establish interbreeding and ancestry in living populations. The manuscript does not give one global percentage because ancestry differs among present

populations and new ancient genomes continue to refine the timing and distribution.

Claims about clothing, shelters and networks are phrased at the level preserved evidence supports. Organic materials decay, so stone and bone overrepresent the surviving toolkit. No private motives, dialogue or undocumented scenes have been added. The disappearance of Neanderthals remains a multi-causal demographic problem, with climate instability, small populations, competition, disease, assimilation and chance carrying different levels of support.

Deglaciation and the Younger Dryas

Clark and colleagues support the onset of Northern Hemisphere deglaciation around 19,000 to 20,000 years ago. Lambeck and colleagues support the large postglacial sea-level rise and its uneven timing. The Bølling-Allerød and Younger Dryas dates use the standard Greenland and North Atlantic event framework, rounded to 14,700 and 12,900 to 11,700 years ago.

Renssen and colleagues tested Younger Dryas mechanisms against proxy evidence and found that weakened Atlantic overturning, moderate negative radiative forcing and altered atmospheric circulation together gave the best fit. No mechanism alone reproduced the full signal. The book therefore rejects both a worldwide second ice age and a single settled trigger. The North Atlantic experienced the strongest cold response; other regions differed, and parts of the Southern Hemisphere did not cool in parallel.

Megafaunal extinction

The extinction section treats causation as geographically and taxonomically variable. Barnosky and colleagues concluded that humans contributed on some continents while hunting alone did not explain every pattern. Sandom and colleagues found a strong global association between extinction severity and human palaeobiogeography, with a much weaker climate association in their macroecological analysis. Cooper and colleagues linked abrupt warming events to Holarctic population turnover using ancient DNA and radiocarbon time series. Lorenzen and colleagues found different combinations for different species.

A recent synthesis by Svenning and colleagues gives strong weight to human expansion and notes that earlier Pleistocene climate oscillations did not produce the same selective global loss of large animals. A 2025 systematic review by Mathew Stewart and colleagues found that scientific opinion remains deeply divided across disciplines and methods. Other regional studies assign more weight to climate in particular settings. The manuscript's most contestable causal judgement is that expanding human pressure mattered strongly and was sometimes decisive while habitat change and demography altered local outcomes. This is firmer than a neutral split and narrower than one universal overkill claim.

Wrangel Island mammoth survival to roughly 4,000 years ago follows ancient DNA, radiocarbon and environmental-DNA evidence summarised by Stuart and Wang and colleagues. It is used to show staggered population extinction, not to date the disappearance of mainland mammoths.

Present climate comparison

The statement that modern warming has a different forcing and pace from orbital glacial cycles follows the physical-science assessment of the Intergovernmental Panel on Climate Change. The Ice Age record demonstrates climate sensitivity, feedback and regional change; it does not make modern industrial forcing an ordinary Milankovitch transition. Current material was rechecked on 4 September 2026.

How we know

The evidential summary draws on Woodward, Alley and the research articles listed below. Landforms can be erased or reused by later ice. Oxygen isotopes combine ice-volume and temperature effects. Ice-core chronologies are strongest where annual layers can be counted and become less direct at depth. Radiocarbon requires calibration and loses power near its practical age limit. Ancient DNA preservation is geographically biased and contamination-sensitive. Archaeology favours durable materials and intensively studied regions.

The broad existence, sequence and scale of repeated glaciation are established by independent records. The weakest material retained relative to memorability is the exact ecological role of mammoths and other herbivores in maintaining open steppe. It is stated as a possible contribution rather than a dominant or universal cause.

Bibliography

Chronology, data and original research

Clark, Peter U., Arthur S. Dyke, Jeremy D. Shakun, Anders E. Carlson, Jorie Clark, Barbara Wohlfarth, Jerry X. Mitrovica, Steven W. Hostetler, and A. Marshall McCabe. "The Last Glacial Maximum." *Science* 325, no. 5941 (2009): 710-714. doi:10.1126/science.1172873.

Cooper, Alan, Chris Turney, Konrad A. Huguen, Barry W. Brook, H. Gregory McDonald, and Corey J. A. Bradshaw. "Abrupt Warming Events Drove Late Pleistocene Holarctic Megafaunal Turnover." *Science* 349, no. 6248 (2015): 602-606. doi:10.1126/science.aac4315.

EPICA Community Members. "Eight Glacial Cycles from an Antarctic Ice Core." *Nature* 429 (2004): 623-628. doi:10.1038/nature02599.

Hays, J. D., John Imbrie, and N. J. Shackleton. "Variations in the Earth's Orbit: Pacemaker of the Ice Ages." *Science* 194, no. 4270 (1976): 1121-1132. doi:10.1126/science.194.4270.1121.

Higham, Tom, Katerina Douka, Rachel Wood, Christopher Bronk Ramsey, et al. "The Timing and Spatiotemporal Patterning of Neanderthal Disappearance." *Nature* 512 (2014): 306-309. doi:10.1038/nature13621.

International Commission on Stratigraphy. *International Chronostratigraphic Chart*. Version 2026/06. 2026.

Lambeck, Kurt, Hélène Rouby, Anthony Purcell, Yiyi Sun, and Malcolm Sambridge. "Sea Level and Global Ice Volumes from the Last Glacial Maximum to the Holocene." *Proceedings of the National Academy of Sciences of the United States of America* 111, no. 43 (2014): 15296-15303. doi:10.1073/pnas.1411762111.

Lisiecki, Lorraine E., and Maureen E. Raymo. "A Pliocene-Pleistocene Stack of 57 Globally Distributed Benthic Oxygen-18 Records." *Paleoceanography* 20 (2005): PA1003. doi:10.1029/2004PA001071.

Lorenzen, Eline D., David Nogués-Bravo, Ludovic Orlando, et al. "Species-Specific Responses of Late Quaternary Megafauna to Climate and Humans." *Nature* 479 (2011): 359-364. doi:10.1038/nature10574.

Lüthi, Dieter, Martine Le Floch, Bernhard Bereiter, et al. "High-Resolution Carbon Dioxide Concentration Record 650,000-800,000 Years Before Present." *Nature* 453 (2008): 379-382. doi:10.1038/nature06949.

Meyer, Matthias, Martin Kircher, Marie-Theres Gansauge, et al. "A High-Coverage Genome Sequence from an Archaic Denisovan Individual." *Science* 338, no. 6104 (2012): 222-226. doi:10.1126/science.1224344.

Prüfer, Kay, Cesare de Filippo, Steffi Grote, et al. "A High-Coverage Neandertal Genome from Vindija Cave in Croatia." *Science* 358, no. 6363 (2017): 655-658. doi:10.1126/science.aao1887.

Renssen, Hans, Aurélien Mairesse, Hugues Goosse, Pierre Mathiot, Oliver Heiri, Didier M. Roche, Kerim H. Nisancioglu, and Paul J. Valdes. "Multiple Causes of the Younger Dryas Cold Period." *Nature Geoscience* 8, no. 12 (2015): 946-949. doi:10.1038/ngeo2557.

Sandom, Christopher J., Søren Faurby, Brody Sandel, and Jens-Christian Svenning. "Global Late Quaternary Megafauna Extinctions Linked to Humans, Not Climate Change." *Proceedings of the Royal Society B: Biological Sciences* 281 (2014): 20133254. doi:10.1098/rspb.2013.3254.

Tierney, Jessica E., Jiang Zhu, Jonathan King, Steven B. Malevich, Gregory J. Hakim, and Christopher J. Poulsen. "Glacial Cooling and Climate Sensitivity Revisited." *Nature* 584 (2020): 569-573. doi:10.1038/s41586-020-2617-x.

Wang, Yucheng, Mikkel Winther Pedersen, Inger Greve Alsos, et al. "Late Quaternary Dynamics of Arctic Biota from Ancient Environmental Genomics." *Nature* 600 (2021): 86-92. doi:10.1038/s41586-021-04016-x.

Reviews and syntheses

Barnosky, Anthony D., Paul L. Koch, Robert S. Feranec, Scott L. Wing, and Alan B. Shabel. "Assessing the Causes of Late Pleistocene Extinctions on the Continents." *Science* 306, no. 5693 (2004): 70-75. doi:10.1126/science.1101476.

Herbert, Timothy D. "The Mid-Pleistocene Climate Transition." *Annual Review of Earth and Planetary Sciences* 51 (2023): 389-418. doi:10.1146/annurev-earth-032320-104209.

Hewitt, Godfrey. "The Genetic Legacy of the Quaternary Ice Ages." *Nature* 405 (2000): 907-913. doi:10.1038/35016000.

Intergovernmental Panel on Climate Change. *Climate Change 2021: The Physical Science Basis*. Edited by Valérie Masson-Delmotte, Panmao Zhai, Anna Pirani, et al. Cambridge: Cambridge University Press, 2021. doi:10.1017/9781009157896.

Stewart, John R., Adrian M. Lister, Ian Barnes, and Love Dalén. "Refugia Revisited: Individualistic Responses of Species in Space and Time." *Proceedings of the Royal Society B: Biological Sciences* 277 (2010): 661-671. doi:10.1098/rspb.2009.1272.

Stewart, Mathew, Carli Peters, Michael J. Ziegler, W. Christopher Carleton, Patrick Roberts, Nicole Boivin, and Huw S. Groucutt. "The State of the Late Quaternary Megafauna Extinction Debate: A Systematic Review and Analysis." *Frontiers in Mammal Science* 4 (2025): 1678231. doi:10.3389/fmamm.2025.1678231.

Svenning, Jens-Christian, Rhys T. Lemoine, Juraj Bergman, Robert Buitenwerf, Elizabeth Le Roux, Erick Lundgren, Ninad Mungi, and Rasmus Ø. Pedersen. "The Late-Quaternary Megafauna Extinctions: Patterns, Causes, Ecological Consequences and Implications for Ecosystem Management in the Anthropocene." *Cambridge Prisms: Extinction* 2 (2024): e5. doi:10.1017/ext.2024.4.

Zachos, James, Mark Pagani, Lisa Sloan, Ellen Thomas, and Katharina Billups. "Trends, Rhythms, and Aberrations in Global Climate 65 Ma to Present." *Science* 292, no. 5517 (2001): 686-693. doi:10.1126/science.1059412.

Books

Alley, Richard B. *The Two-Mile Time Machine: Ice Cores, Abrupt Climate Change, and Our Future*. Princeton: Princeton University Press, 2000.

Gaffney, Vincent, Simon Fitch, and David Smith. *Europe's Lost World: The Rediscovery of Doggerland*. CBA Research Report 160. York: Council for British Archaeology, 2009.

Stuart, Anthony J. *Vanished Giants: The Lost World of the Ice Age*. Chicago: University of Chicago Press, 2021.

Woodward, Jamie. *The Ice Age: A Very Short Introduction*. Oxford: Oxford University Press, 2014.

Wragg Sykes, Rebecca. *Kindred: Neanderthal Life, Love, Death and Art*. London: Bloomsbury Sigma, 2020.